

GIS-BASED ASSESSMENT OF THE LANDFORM DISTRIBUTION OF 2100 PREDICTED CLIMATE CHANGE AND ITS INFLUENCE ON BIODIVERSITY AND NATURAL PROTECTED AREAS IN ROMANIA

PETRIȘOR Alexandru Ionuț

Abstract. Potential effects of climate change include loss of biodiversity and ecodiversity due to direct influences and habitat changes and impacts on agricultural systems resulting into the decrease of productivity. While direct assessment is rarely possible, the evaluation methodology focused on statistical modelling. This study used the Geographical Information Systems to look at the spatial distribution of 2100 predicted temperatures in Romania by landform and biogeographical regions, assess their potential influences on natural protected areas, and model the shift of biogeographical regions under the hypothesis according to which increased temperature result into the expansion of warmer regions over the colder ones. Results suggest that, according to the predictions, the most important threats occur in natural protected areas from the mountain regions and in agriculture, and that the Steppic region could expand over the Continental and Black Sea ones, with caveats due to the hypothesis and methodology.

Keywords: biodiversity, ecodiversity, climate change, biogeographical region.

Rezumat. Evaluarea în Sistem Informațional Geografic a distribuției schimbărilor climatice prezise pentru 2100 în funcție de relief și a influenței acesteia asupra biodiversității și ariilor naturale protejate din România. Efectele potențiale ale schimbărilor climatice includ erodarea biodiversității și ecodiversității datorită influenței directe și modificării habitatelor și impacturi asupra agrosistemelor, care conduc la scăderea productivității. Deși evaluarea directă este arareori posibilă, metodologia de evaluare s-a concentrat asupra modelării statistice. Acest studiu a folosit Sistemele Informaționale Geografice pentru a analiza distribuția spațială a predicțiilor pe 2100 a temperaturii din România în funcție de relief și regiunea biogeografică, pentru a evalua influența potențială asupra ariilor naturale protejate și a modela translația regiunilor biogeografice sub ipoteza potrivit căreia creșterea temperaturii determină extinderea regiunilor calde peste cele reci. Rezultatele arată că, pe baza predicțiilor, cele mai importante amenințări apar în ariile naturale protejate din regiunile montane și în agricultură și că regiunea de stepă s-ar putea extinde peste cea continentală și peste cea a Mării Negre, sub rezerva ipotezei și a metodologiei.

Cuvinte cheie: biodiversitate, ecodiversitate, schimbări climatice, regiune biogeografică.

INTRODUCTION

Biological diversity means the variability among living organisms from all sources including, *inter alia*, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystems (UNITED NATIONS, 1993); the latter is called ecodiversity. Based on the spatial scale, several levels of diversity are defined: α - diversity of ecosystems, communities, biocoenoses, taxonomic or functional groups; β - diversity of ecosystems within a complex of ecosystems, diversity of habitats or gradients; γ - diversity of regional complexes of ecosystems, *e.g.* the European biogeographical regions; δ - diversity of macro-regional complexes of ecosystems, *e.g.* the global biogeographical regions; ϵ - diversity of life environments; and ω - phylogenetic diversity (MAGURRAN, 1988; PUSCEDDU, 2008).

Previous studies have indicated that climate change influences all biological and ecological systems, regardless of being natural or man-dominated, protected or not. Species have been affected directly or due to habitat changes (CONDÉ & RICHARD, 2008) with respect to their interactions (MARSHALL et al., 2008), distribution, extinction rates, reproduction timings, length of growing seasons (SECRETARIAT OF THE CONVENTION ON BIOLOGICAL DIVERSITY, 2007). Moreover, most of the important elements of global change have been shown likely to increase the prevalence of biological invaders (DUKES & MOONEY, 1999; DUKES, 2003). Ecosystems have been affected in ways currently difficult to predict due to the influence of climate change on the transport vectors (MARSHALL et al., 2008), due to the exacerbation of the effects of habitat fragmentation (THOMAS, 2003), or due to the influence of the large-scale and regional circulation on local processes (BLENCKNER & CHEN, 2003). Significant changes have been expected in the Alpine ecosystems (CONDÉ & RICHARD, 2008) due to the retreat or disappearance of Alpine species caused by climate change (SECRETARIAT OF THE CONVENTION ON BIOLOGICAL DIVERSITY, 2007). The overall trends regarding glaciers, plants, insects, and temperatures have shown remarkable internal consistency at high elevations (EPSTEIN et al., 2008). Impacts on agricultural ecosystems have included increased exposure to heat stress, changes in rainfall patterns, greater leaching of nutrients, more wildfires, greater erosion due to stronger winds, and spread of pests and diseases (SECRETARIAT OF THE CONVENTION ON BIOLOGICAL DIVERSITY, 2007).

The methodology used to assess the correlation between climate change and loss of biodiversity has included statistical analyses of correlations between climatic and biological data (BLENCKNER & CHEN, 2003), small-scale experiments that study species dynamics under altered precipitation regimes or simulated warming, identification of functional traits that are related to tolerance of different climates, long-term observations of species composition changes correlated with climate variation (DUKES & MOONEY, 1999), and climate modeling using different software (DUKES & MOONEY, 1999; MALCOLM, 2003; SCHRÖTER et al., 2003; EPSTEIN et al., 2008; MARSHALL et al., 2008).

However, while statistical analyses have been suitable for large scale systems and climate models have been designed for analyses at continental scale, intermediate scale analyses have been performed using the Geographical Information Systems (GIS), representing decision support systems involving the integration of spatially referenced data in a problem solving environment (COWEN, 1988). The use of GIS in relationship with biodiversity has been especially productive when looking at β or γ diversity (PETRIȘOR, 2008).

A special inference was needed to analyze the possible expansion of biogeographical regions due to climate change. The underlying hypothesis is that when regions with colder climates and regions with warmer climates are neighboring, the temperature increase in the colder areas results into a gradual colonization of these areas by species migrating from the warmer climate and finally into a replacement of colder biogeographical regions by the warmer ones, given the modification of abiotic conditions as a result of climate change, and of the species composition as a result of migration. The start point in drawing the new boundaries was represented by current temperature ranges characteristic to each biogeographical region, as indicated by different sources: the EBONE / European Biodiversity Observation Network describes average temperatures for the Pannonian region (WAGENINGEN-UR, 2100a), Alpine region (WAGENINGEN-UR, 2100b), and Continental region (WAGENINGEN-UR, 2100c); other sources published data on the Steppic region (SUNDSETH, 2009) and Black Sea region (ZAITSEV et al., 2002). The intervals are displayed in Table 1; as it can easily be noticed, most intervals overlap. The actual overlap is depicted in Fig. 1.

Table 1. Displaying the range of current temperatures characteristic to all Romanian biogeographical regions.
Tabel 1. Intervale de temperatură caracteristice tuturor regiunilor biogeografice din România.

Biogeographical region	Minimum temperature (°C)	Maximum temperature (°C)
Pannonian	-3	27
Alpine	-7	20
Continental	-5	23
Steppic	-15	30
Black Sea	6	25




Figure 1. Displaying the overlapping ranges of current temperatures characteristic to each biogeographical region.
Figura 1. Intervale suprapuse ale temperaturilor caracteristice fiecărei regiuni biogeografice.

The problem due to overlapping does not allow the application of discriminatory analysis that could yield a logical separation of biogeographical regions based solely on temperature ranges. However, there is only one supported inference given the hypothesis and current ranges: minimum predicted temperatures below -7°C and/or maximum predicted temperatures over 27°C characterize the Steppic region.

This study aimed to look at the spatial distribution of predicted climate changes by landform, to evaluate their potential influence on the biodiversity and natural protected areas in Romania, and to predict the spatial shifts of biogeographical regions based on current temperature ranges and the assumption that temperature increases will result into the expansion of warmer regions over the colder ones.

MATERIALS AND METHODS

This study integrated five datasets, summarized in Table 2:

(1 and 2) Climate data consist of global current (1) and predicted (2) temperatures. In addition, we computed the difference between actual and predicted average temperatures for each raster cell. The analysis of differences indicated that within the Romanian territories all values were positive, ranging between 2.329 and 2.7125°C . Consequently, we defined three classes to describe the range of temperature differences: low (2.329 - 2.457°C), average (2.457 - 2.585°C), and high (2.585 - 2.713°C).

(3) Landforms: the main methodological issue was related to the definition of landforms by altitude. Different authors propose a wide range of altitude limits to discriminate between landforms: plains - up to 300 m., hills and plateaus - up to 800 m. (MĂRA, 2007); plains - up to 200 m., hills and plateaus - up to 1000 m. (CAZAN et al., 2004). However, even the two works cited above do not provide a clear limit of separation between hills or plateaus and mountains. Moreover, the two classes overlap for an interval of approximately 200 m altitude. This study utilizes the

following limits of the landforms: floodplain - 0 to less than 20 m, plain - 0 to less than 200 m, hill or plateau - 200 to less than 900 m, and mountain - over 900 m (PETRIȘOR, 2009b).

(4) Data on land cover and use: we used the first level of this classification, defining five land cover classes: artificial surfaces, agricultural areas, forests or semi-natural areas, wetlands, and water bodies (Commission of the European Communities, 1995).

(5) Biogeographical regions: five of the twelve regions identified in Europe are present in Romania: Continental, Steppic, Alpine, Pannonian, and Black Sea.

(6) Data on natural protected areas from Romania contain information on the following types: (a) scientific reserves, natural reserves and natural monuments, (b) national parks and natural parks (biological), (c) Sites of Community Importance, (d) Special Areas of Conservation, (e) Special Protection Areas, and (f) area where the Convention on the Protection and Sustainable Development of the Carpathians is applied, even though the latter cannot be considered a natural protected area in the true meaning of this concept, as protective actions are only recommended, but not compulsory within its perimeter. Categories (c), (d), and (e) were established through the Natura 2000 Programme of the European Union.

Table 2. Specifications on the data used in the study: dataset, provider, location, format, remarks and transformations.
Tabel 2. Specificații asupra datelor utilizate în acest studiu: set de date, sursă, adresă, format, observații și transformări.

No.	Dataset	Provider	URL	Format	Remarks	Transformations
1	Climate - actual	University of Berkeley	http://biogeo.berkeley.edu/worldclim/diva/diva_worldclim_2-5m.zip	DIVA-GIS software (Hijmans <i>et al.</i> , 2001)	Produced by the project WorldClim (Hijmans <i>et al.</i> , 2005); 2.5 min × 2.5 min	Imported in ArcView GIS 3.X, projected into Stereo 1970, subsample for Romania
2	Climate - predicted	University of Berkeley	http://biogeo.berkeley.edu/worldclim/diva/diva_wc_ccm3_2-5m.zip	DIVA-GIS software (Hijmans <i>et al.</i> , 2001)	Predictions for 2100 based on 2×CO ₂ concentration and CCM3 model (GOVINDASAMY <i>et al.</i> , 2003); 2.5 min × 2.5 min	Imported in ArcView GIS 3.X, projected into Stereo 1970, subsample for Romania
3	Land-form	Consultative Group on International Agricultural Research - Consortium for Spatial Information	http://srtm.csi.cgiar.org/SELECTION/inputCoord.asp	Digital Elevation Model (DEM)	Nearly 90 m × 90 m	Import into Arc GIS, then export to ArcView GIS 3.X, projected into Stereo 1970, subsample for Romania
4	Land cover and use data	CORINE (Coordinated Information on the European Environment) Land Cover 2000 (CLC2000) seamless vector database	http://dataservice.eea.europa.eu/dataservice/metadata.asp?id=950	ArcView GIS 3.X	2000 data	Projected into Stereo 1970, subsample for Romania
5	Biogeographical regions	European Environment Agency data services	http://dataservice.eea.europa.eu/dataservice/metadata.asp?id=308	ArcView GIS 3.X	2001 data	Projected into Stereo 1970, subsample for Romania
6	Natural protected areas	Romanian Ministry of the Environment and Sustainable Development	http://www.mmediu.ro/departament_ape/biodiversitate/	ArcView GIS 3.X	Not all types of protected areas legally defined are available	No transformation needed

The datasets were overlaid for a visual and quantitative analysis of information. Quantitative analyses consisted of using GIS spatial analysis and geo-processing functions to compute areas of specific classes describing the overlapping of the three temperature intervals and either landforms, biodiversity, or natural protected areas. In addition, we computed:

(a) the percentage represented by each combination between the temperature interval and class of specific feature from all temperature intervals, to pinpoint which temperature intervals have more influence on specific feature classes, and

(b) the percentage represented by each combination between the temperature interval and class of specific feature from all feature classes, to analyze which feature classes are more impacted by a certain temperature interval.

To analyze the expansion of regions given the increases of temperatures and provided the methodological limitation described in Figure 1 and Table 1, GIS was used to determine the areas where the minimum predicted temperatures are below -7^oC and/or the maximum predicted temperatures are over 27^oC. However, the results indicated that such areas are situated in the Continental, Pannonian, and Black Sea regions, where maximum predicted temperatures exceed current minimum values, and in the Alpine region, where minimum temperatures fall below current minimum values. While colonization of the Black Sea and Continental regions could be possible due to spatial proximity, the results were modified to eliminate areas situated in the Alpine and Pannonian regions due to the lack of spatial continuity and impossibility of colonization.

RESULTS AND DISCUSSION

First of all, it is important to reiterate that the differences between the predicted and actual temperatures are positive at all locations composing the territory of Romania, ranging between 2.329 and 2.7125.

The results are presented in Figures 2 - 7. Figure 2 displays the distribution of low, average, and respectively high differences between actual and predicted average temperature by landforms, land cover classes, biogeographical regions, and types of natural protected area. Figure 3 displays information from a reverted perspective, looking at the influence of the classes of differences between actual and predicted average temperature on each type of landform, land cover class, biogeographical region, and type of natural protected area. Figure 4 displays the results of the analyses of the spatial distribution of low, average, and high differences between the predicted and actual temperature based on the landform, Figure 5 displays the spatial relationship between climate predictions and biodiversity, assessed by using the first level of CORINE land cover, and Figure 6 displays the potential impact of increased average temperatures on existing natural protected areas. Figure 7 redraws the boundary of biogeographical regions based on the comparison between predictions and current ranges of temperatures.

This study aimed to analyze the spatial distribution of predicted climate changes by landform and evaluate their potential impact on the biodiversity and natural protected areas in Romania.

Our findings suggest that predicted climate changes, reflected by positive differences between the predicted and actual average temperature, are not distributed uniformly over the Romanian landforms (Figs. 2-4). The highest temperature differences occurred mostly in the plain areas (64.5%), average differences in the floodplains (72.86%), and low differences characterized floodplains (52.26%) and hills or plateaus (44.13%). With respect to the landform, floodplains were dominated by average differences (61.99%), plains by high differences (87.22%), hills or plateaus by low differences (56.5%), and mountains by high differences (98.1%).

The impact on biodiversity can be assessed by looking at the relationship between predicted temperature differences and diversity assessed using the land cover and biogeographical regions (Figs. 2, 3, and 5). In the first case, high differences affected the agricultural surfaces and forest or semi-natural areas (47.15, respectively 44.92%), while both average and low differences impacted only the agricultural surfaces (74.12, respectively 63.24%). With respect to the land cover classes, artificial areas received predominantly high temperatures (65.88%), agricultural areas were impacted by high (50.67%) and average (40.1%) temperature differences, forest or semi-natural areas were affected by high differences (80.98%), and wetlands and water bodies by low differences (63.25%, respectively 45.95%). In the second case, high and average differences had a significant impact mostly on the Continental region (61.67%, respectively 58.64%), while low differences predominated in the Steppic region (81.94%). With respect to the biogeographical region, high differences were expected in the Continental and Alpine regions (67.63, respectively 99.82%), the Pannonian region was characterized by average differences (96.47%), and the Black Sea region was situated exclusively within the low differences range (100%). The Steppic region was mostly affected by average and low differences (42.45, respectively 41.70%).

Finally, the analysis of the potential impact of predicted temperature differences on the natural protected areas, summarized in Figures 2, 3, and 6, indicated that high and average temperature differences affected the area where the Convention on the Protection and Sustainable Development of the Carpathians is applied (58.66%, respectively 31.84%), while low temperature differences were predominant in the SPAs. High temperature differences were the most common regardless of the type of natural protected area (scientific and natural reserves and natural monuments - 62.82%, national and natural parks - 48.95%, SCIs - 59.51%, the area under the Convention on the Protection of the Carpathians - 88.68%, SPAs - 49.37%, and SACs - 73.37%).

Overall, the evolution of average temperatures in Romania suggests an increasing trend, with positive differences between the current and predicted values all over the national territory, ranging between 2.329 and 2.7125. Our results, consistent with the findings of CONDÉ & RICHARD (2008) and the SECRETARIAT OF THE CONVENTION ON BIOLOGICAL DIVERSITY (2007), also indicate that the highest temperature differences will occur in the mountain areas (Fig. 4), situated in the Alpine biogeographical region (Fig. 5) and covered in majority by forests or semi-natural areas and agricultural surfaces. Since most of the Romanian natural protected areas are situated at high altitudes (the percentage ranged from 55% for SACs to 93% for SPAs, excluding the area where the Convention on the Protection and Sustainable Development of the Carpathians is applied, situated exclusively at high altitudes - PETRIȘOR, 2009b), the additional pressure due to climate changes will significantly threaten the existing natural protected areas.

Moreover, since most of the Romanian territory is covered by agricultural surfaces, the influence of high temperatures on floodplains and plains, and on agricultural surfaces (Figs. 1 and 2) will most likely result into a decrease of the agricultural production (consistent with the results published by the SECRETARIAT OF THE CONVENTION ON BIOLOGICAL DIVERSITY, 2007) and would potentially affect the entire economy.

Figure 7 indicates a possible expansion of the Steppic region over the Black Sea region and over some parts of the Continental region. Moreover, the predicted maximum temperatures in the current Steppic region exceed current maximum values, which could result into disappearance of species less tolerant to high temperatures. However, these findings are subject to caveats due to the hypothesis and due to the methodological limitation of discriminating between the intervals. Another limitation is that the hypothesized process will have a time span characteristic to the large regional complexes of ecosystems, exceeding the prediction horizon, excepting perhaps for pioneer and invasive species.




Figure 2. Distribution of low, average, and respectively high differences between actual and predicted average temperature by landforms, land cover classes, biogeographical regions, and types of natural protected area - scientific and natural reserves and natural monuments (R), national and natural parks (P), Sites of Community Importance (SCI), area under the Convention on the Protection of the Carpathians (C), Special Protection Areas (SPA), and Special Areas of Conservation (SAC). Temperature differences are grouped in three classes: low (2.329-2.457), average (2.457-2.585), and high (2.585-2.713).

Figura 2. Distribuția diferențelor mici, medii și mari dintre temperaturile existente și prezisă în funcție de relief, categoriile de acoperire a terenului, regiunile biogeografice și tipurile de arii naturale protejate - rezervații științifice și naturale și monumente naturale (R), parcuri naționale și naturale (P), situri de importanță comunitară (SCI), zona de aplicare a Convenției Carpatice (C), arii de protecție avifaunistică (SPA), și arii speciale de conservare (SAC). Diferențele de temperatură sunt grupate în trei categorii: mici (2,329-2,457), medii (2,457-2,585) și mari (2,585-2,713).

CONCLUSIONS

In summary, findings indicate that predicted climate change for 2100 could possibly affect the biodiversity of Romania. Most important changes will occur in the mountain regions that are already a priority on the European environmental agenda. Other changes will occur in the continental region, which could result into the transformation of some parts of it into a steppe. However, the later statement should be interpreted with the caveats of the hypothesis according to which temperature increase lead to the expansion of warmer regions over the colder ones, and the methodological limitations. Last but not least, the changes are likely to affect natural protected areas, especially those situated at higher altitudes.

ACKNOWLEDGEMENTS

The author would like to thank PhD. Sorin Cheval from the National Meteorological Administration for his valuable comments and suggestions that helped improving the study design, as well as the elaboration of this article.




Figure 3. Influence of the classes of differences between actual and predicted average temperature on each type of landform, land cover class, biogeographical region, and type of natural protected area - scientific and natural reserves and natural monuments (R), national and natural parks (P), Sites of Community Importance (SCI), area under the Convention on the Protection of the Carpathians (C), Special Protection Areas (SPA), and Special Areas of Conservation (SAC). Temperature differences are grouped in three classes: low (2.329-2.457), average (2.457-2.585), and high (2.585-2.713).

Figure 3. Influența claselor de diferențe dintre temperaturile existentă și prezisă asupra fiecărei unități de relief, fiecărei categorii de acoperire a terenului, fiecărei regiuni biogeografice și fiecărui tip de arii naturale protejate - rezervații științifice și naturale și monumente naturale (R), parcuri naționale și naturale (P), situri de importanță comunitară (SCI), zona de aplicare a Convenției Carpatice (C), arii de protecție avifaunistică (SPA), și arii speciale de conservare (SAC). Diferențele de temperatură sunt grupate în trei categorii: mici (2,329-2,457), medii (2,457-2,585) și mari (2,585-2,713).




Figure 4. Distribution of 2100 predicted temperature increase by landform in Romania. The magnitude of temperature differences is displayed using gray shades; darker shades indicate higher values, suggesting an important increase of the average temperature.

Figura 4. Distribuția temperaturilor prezise pentru 2100 în funcție de relief în România. Mărimea diferențelor este indicată prin nuanțe de gri; nuanțele închise indică valori mari, sugerând creșteri importante ale temperaturii medii.




Figure 5. Distribution of 2100 predicted temperature increase in the Romanian biogeographical regions. The magnitude of temperature differences is displayed using gray shades; darker shades indicate higher values, suggesting a significant increase of the average temperature.

Figura 5. Distribuția temperaturilor prezise pentru 2100 în regiunile biogeografice din România. Mărimea diferențelor este indicată prin nuanțe de gri; nuanțele închise indică valori mari, sugerând creșteri importante ale temperaturii medii.




Figure 6. Distribution of 2100 predicted temperature increase in the Romanian natural protected areas. The magnitude of temperature differences is displayed using gray shades; darker shades indicate higher values, suggesting a significant increase of the average temperature.

Figura 6. Distribuția temperaturilor prezise pentru 2100 în ariile naturale protejate din România. Mărimea diferențelor este indicată prin nuanțe de gri; nuanțele închise indică valori mari, sugerând creșteri importante ale temperaturii medii.




Figure 7. 2100 predicted shift of biogeographical regions. The image displays the limits of the expanded Steppic region; its current boundaries correspond to an area where the predictions indicate that current maximum temperatures will be exceeded, possibly resulting into the disappearance of some species and ecosystems.

Figura 7. Translații ale regiunilor biogeografice prezise pentru 2100. Imaginea arată limita de extindere a regiunii de stepă; limitele actuale corespund unei regiuni unde conform predicțiilor vor fi depășite temperaturile maxime actuale, ceea ce probabil va conduce la dispariția unor specii și ecosisteme.

REFERENCES

- BLENCKNER T. & CHEN D. 2003. *Comparison of the impact of regional and North Atlantic atmospheric circulation on an aquatic ecosystem*. *Climate Research*. **23**: 131-136.
- CAZAN C. M., CHERCIU R., MANDA M. 2004. *Creează-ți mediul! Dicționar de termeni*. Holcim. Bucharest: 158 pp.
- CONDÉ S. & RICHARD D. 2008. *Europe's biodiversity - biogeographical regions and seas*. European Environment Agency. Copenhagen: 3-4.
- COWEN D. J. 1988. *GIS versus CAD versus DBMS: What Are the Differences?* *Photogrammetric Engineering & Remote Sensing*. **54**: 1551-1555.
- DUKES J. 2003. *Hotter and weedier? Effects of climate change on the success of invasive species*. In: GREEN R. E., HARLEY M., MILES J., SCHARLEMANN J., WATKINSON A., WATTS O. *Global Climate Change and Biodiversity*. University of East Anglia. Norwich: 25.
- DUKES J. S. & MOONEY H. A. 1999. *Does global change increase the success of biological invaders?* *Trends in Ecology and Evolution*. **14**(4): 135-139.
- EPSTEIN P. R., DIAZ H. F., ELIAS S., GRABHERR G., GRAHAM N. E., MARTENS W. J. M., MOSLEY-THOMPSON E., SUSSKIND J. 1998. *Biological and Physical Signs of Climate Change: Focus on Mosquito-borne Diseases*. *Bulletin of the American Meteorological Society*. **79**(3): 409-417.
- GOVINDASAMY B., DUFFY P. B., COQUARD J. 2003. *High-resolution simulations of global climate, Part 2: Effects of increased greenhouse gases*. *Climate Dynamics*. **21**: 391-404.
- HIJMANS R. J., GUARINO L., CRUZ M., ROJAS E. 2001. *Computer tools for spatial analysis of plant genetic resources data: 1. DIVA-GIS*. *Plant Genetic Resources Newsletter*. **127**: 15-19.
- HIJMANS, R. J., CAMERON S. E., PARRA J. L., JONES P. G., JARVIS A. 2005. *Very high resolution interpolated climate surfaces for global land areas*. *International Journal of Climatology*. **25**: 1965-1978.
- MAGURRAN A. E. 1988. *Ecological Diversity and Its Measurement*. Princeton University Press. Princeton: 81-99.
- MALCOLM J. R. 2003. *Models of the global impact of climate change on biodiversity and adaptive responses*. In: GREEN R. E., HARLEY M., MILES J., SCHARLEMANN J., WATKINSON A., WATTS O. *Global Climate Change and Biodiversity*. University of East Anglia. Norwich: 6-7.
- MARSHALL J. D., BLAIR J. M., PETERS D. P. C., OKIN G., RANGO A., WILLIAMS M. 2008. *Predicting and understanding ecosystem responses to climate change at Continental scales*. *Frontiers in Ecology and the Environment*. **6**(5): 273-280.
- MÂRA L. 2007. *Elaborarea studiului de analiză strategică de mediu pentru Planul de Amenajare a Teritoriului Național, Secțiunea a VI-a - Zone turistice. Faza 1 - Aspecte relevante ale stării actuale a mediului, caracteristici de mediu, Zone cu importanță specială pentru mediu*. IPTANA-S.A. Bucharest: 235 pp.
- PETRIȘOR A. I. 2008. *Levels of biological diversity: a spatial approach to assessment methods*. *Romanian Review of Regional Studies*. Cluj-Napoca. **4**(1): 41-62.
- PETRIȘOR A. I. 2009a. *Spatial approach to the assessment of anthropic impact on biodiversity based on the Nomenclature of Territorial Units for Statistics (NUTS) applicable to Romania*. *Oltenia. Studies and communications*. *Nature Sciences*. **25**: 305-308.
- PETRIȘOR A. I. 2009b. *GIS assessment of geodiversity covered by natural protected areas in Romania*. *Studia Universitatis Vasile Goldiș. Life Sciences Series*. **19**(2): 359-363.
- PUSCEDDU A. 2008. *Biodiversità*. In: *Fondamenti di Analisi dei Sistemi Ecologici*. Università Politecnica delle Marche. Ancona: 6-7.
- SCHRÖTER D., ZAEHLE S., SCHAPHOFF S., LUCHT W., CRAMER W. 2003. *Modeling global change effects on vegetation and exploring our vulnerability*. In: GREEN R. E., HARLEY M., MILES J., SCHARLEMANN J., WATKINSON A., WATTS O. *Global Climate Change and Biodiversity*. University of East Anglia. Norwich: 4-5.
- SUNDSETH K. 2009. *Natura 2000 in the Steppic Region*. Office for Official Publications of the European Communities. Luxembourg: 8 pp.
- THOMAS C. 2003. *Modeling global change effects on vegetation and exploring our vulnerability*. In: GREEN R. E., HARLEY M., MILES J., SCHARLEMANN J., WATKINSON A., WATTS O. *Global Climate Change and Biodiversity*. University of East Anglia. Norwich: 22-23.
- WAGENINGEN-UR. 2100a. *Pannonic-Pontic (PAN), European Biodiversity Observation Network*. Available at the URL <http://www.ebone.wur.nl/UK/Project+background/European+Environmental+Stratification/Pannonic-Pontic+%28PAN%29/>.
- WAGENINGEN-UR. 2100b. *Alpine South (ALS), European Biodiversity Observation Network*. Available at the URL <http://www.ebone.wur.nl/UK/Project+background/European+Environmental+Stratification/Alpine+South+%28ALS%29/>.
- WAGENINGEN-UR. 2100c. *Continental (CON), European Biodiversity Observation Network*. Available at the URL <http://www.ebone.wur.nl/UK/Project+background/European+Environmental+Stratification/Continental+%28CON%29/>.

- ZAITSEV Y. P., ALEXANDROV B. G., BERLINSKY N. A., ZENETOS A. 2002. *Europe's biodiversity - biogeographical regions and seas. Seas around Europe. The Black Sea - an oxygen-poor sea*. European Environment Agency. available at the URL http://www.eea.europa.eu/publications/report_2002_0524_154909/BlackSea.pdf.
- ***. Commission of The European Communities. 1995. *CORINE land cover: Part One - Methodology*. European Environment Agency. Copenhagen: 163 pp.
- ***. Secretariat of The Convention on Biological Diversity. 2007. *Biodiversity and Climate Change. International Day for Biological Diversity Booklet*. Secretariat of the Convention on Biological Diversity. Montreal: 48 pp.
- ***. United Nations. 1993. *Multilateral Convention on biological diversity (with annexes)*. Concluded at Rio de Janeiro on 5 June 1992. United Nations - Treaty Series 1760(30619): 142-382.

Petrișor Alexandru-Ionuț

Department of Urban and Landscape Planning, School of Urbanism and Landscape Architecture
"Ion Mincu" University of Architecture and Urbanism, Bucharest
Str. Academiei. no. 18-20. sector 1. cod 010014. Bucharest. Romania
Department of Spatial Planning, National Institute for Research and Development in Constructions,
Urbanism and Sustainable Spatial Development URBAN-INCERC, URBANPROIECT branch, Bucharest
Str. Nicolae Filipescu. no. 53-55. sector 2. cod 020961. Bucharest. Romania
E-mail: a.i.petrisor@gmail.com,

Received: January 6, 2010
Accepted: July 15, 2010