

NEW MAASTRICHTIAN MICROVERTEBRATES FROM THE RUSCA MONTANĂ BASIN (ROMANIA)

VASILE Ștefan, CSIKI Zoltán

Abstract. Known since the beginning of the 20th century for their abundant and diverse paleofloristic content, the Maastrichtian deposits of the Rusca Montană Basin yielded no vertebrate fossils until recently, when sauropod, ornithopod, theropod, chelonian, crocodylian and multituberculate remains have been discovered in floodplain fines and river channel sandstones cropping out in the eastern part of the basin. This paper presents additional fossils from a potentially new site within the same area, belonging to taxa and metataxa previously unknown in the Rusca Montană Basin. The fossil assemblage includes anurans, albanerpetontids, lizards, ziphodont crocodylians, additional small theropods, eggshells, freshwater gastropods and angiosperm fructifications. The faunal and sedimentological data suggest that the fossil assemblage is largely autochthonous, and thus reflects the composition of a wetland, poorly drained distal floodplain paleo-ecosystem.

Keywords: microvertebrates, Maastrichtian, Rusca Montană Basin.

Rezumat. Noi microvertebrate maastrichtiene din Bazinul Rusca Montană (România). Cunoscute încă de la începutul secolului al 20-lea datorită unui conținut paleofloristic bogat și divers, depozitele maastrichtiene ale Bazinului Rusca Montană nu au furnizat vertebrate fosile până recent, când resturi ale unor sauropode, ornithopode, theropode, chelonieni, crocodilienii și multituberculate au fost descoperite în depozite fine de câmpie inundabilă și gresii de canal fluvial care aflorază în partea de est a bazinului. Această lucrare prezintă o asociație de fosile provenind dintr-un sit potențial nou din aceeași zonă, fosile ce reprezintă taxoni și metataxonii anterior necunoscuți din Bazinul Rusca Montană. Asociația fosilă include anure, albanerpetontide, crocodili ziphodonți, șopârle, noi taxoni de theropode, coji de ouă, gastropode de apă dulce și fructificații de angiosperme. Datele faunistice și de sedimentologie sugerează o asociație fosilă autohtonă, ce reflectă compoziția unui paleo-ecosistem umed, de câmpie inundabilă distală, slab drenată.

Cuvinte cheie: microvertebrate, Maastrichtian, Bazinul Rusca Montană.

INTRODUCTION

Placed in the southwestern part of the Poiana Ruscă Mountains, the Rusca Montană Basin represents an intramontainous depression bounded by the Bistra Valley to the south and the Neogene Caransebeș Basin to the southwest. The basin formed after the Austrian and "Laramidian" tectogeneses, as a depression lying between the Getic and Supragetic Nappes, both of which cover the Danubian Units (STRUTINSKI & HANN, 1986).

The oldest sedimentary units covering the metamorphic basement are Lower Jurassic marine conglomerates, followed by Middle and Upper Jurassic reef limestones (CANTUNIARI, 1937; DINCĂ, 1977; MAMULEA, 1955). During the Cretaceous, sea level changes led to the deposition of alternating marine and continental sediments: Albian? bauxites, followed by a marine succession with Cenomanian conglomerates and sandstones, Turonian-Coniacian sandy marls and marls, topped by Santonian-Campanian flysch deposits, and finally Maastrichtian continental, detritic and volcanoclastic deposits (CANTUNIARI, 1937; DINCĂ, 1977; MAMULEA, 1955). The sedimentary succession reaches a little more than 2,000 m in thickness, much of which is represented by the Maastrichtian molasse deposits.

Although no formal lithostratigraphic units have yet been described from the Rusca Montană Basin, the Maastrichtian deposits appear under two distinct lithologies. Coarse deposits represented by red, purple, and sometimes grey conglomerates and sandstones occur in the western part of the basin. These are made up of metamorphic and limestone clasts reworked from the underlying basement or older sedimentary deposits, and are bonded in calcareous cementum (DINCĂ, 1977; DUȘA, 1974, 1987; GRIGORESCU, 1992). Mixed tuffaceous and fine-grained detritic deposits occur in the eastern part of the basin, being crossed by Banatitic volcanic intrusions and hosting a few coal lenses (BIȚOIANU, 1970; DINCĂ, 1977; DUȘA, 1970, 1974, 1987; DUȘA & BĂRILĂ, 1973; GRIGORESCU, 1992; NOPCSA, 1905; MAMULEA, 1955). The Maastrichtian age of the later deposits has been based on their rich fossil macroflora (GIVULESCU, 1966, 1968; PETRESCU & DUȘA, 1970, 1980, 1985; SCHAFARZIK, 1906; TUZSON, 1913) and palynological content (ANTONESCU et al., 1983). Despite their similarity in lithology, flora and palynomorphs to contemporaneous deposits from the neighbouring Hațeg Basin (ANTONESCU et al., 1983; DINCĂ et al., 1972; GRIGORESCU, 1992; NOPCSA, 1905), no vertebrate remains have been discovered in over one century of geological research in the Maastrichtian of the Rusca Montană Basin.

The first vertebrates have been reported only recently (CODREA et al., 2009, in press; FEIGI et al., 2010) from the eastern part of the basin, near Lunca Cernii de Sus, from red fine-grained floodplain mudstones and coarser, gray, river channel sandstones. The specimens described are taxonomically diverse, belonging to sauropods (a probable titanosaur), ornithopods, theropods (indeterminate velociraptorines), crocodylians, chelonians, and multituberculate mammals, an assemblage reminiscent of that already known from the Maastrichtian deposits of the Hațeg Basin (e.g. GRIGORESCU, 2010: Table 1). Besides these micro- and macrovertebrate remains, the presence of minute plant seeds and amber-like grains was also mentioned.

MATERIAL AND METHODS

Several outcropping red and gray-blue coloured fine-grained beds have been identified during a field survey along Fărcădeana Brook, a small left-side tributary of Negoiu Creek, within the inhabited area of Negoiu village, Hunedoara County (Fig. 1). About 70 kg of matrix has been sampled from three such beds (one red and two gray-blue in colour) to test for possible microvertebrate remains. The sampled overbank deposits are discontinuous and occur at various heights on the slopes from the brook base, which is gently dipping approximately from northwest to southeast. Since the different beds cannot be followed along the valley, the construction of a synthetic geological log is extremely difficult. The samples have been screen-washed using 2 mm and 0.71 mm mesh size sieves, following the procedure of sediment processing first used for Cretaceous microvertebrates in Romania by GRIGORESCU et al. (1985). The resulting material was sorted under a Zeiss-Stemi binocular microscope. The sample collected from the red-coloured mudstone was not fossiliferous; only a few gastropod opercula were found in one of the gray-blue mudstone samples, but the other yielded numerous small gastropods (opercula and shells), fruits, and microvertebrate remains, a remarkable assemblage described herein. The most representative specimens have been photographed using a Cannon EOS 1000D digital camera mounted on a Zeiss Stemi 2000-C microscope.

RESULTS AND DISCUSSIONS

The microvertebrate-bearing bed that produced the fossil remains consists of about 2 m of massive gray-blue muscovite-rich mudstone, slightly coarser, sandy, in its upper third, being unconformably covered by a well-sorted conglomerate, composed of sub-rounded, mainly quartzitic grains (2.5-5 cm in diameter), along with rare, similarly-sized micaschist clasts. Minute pyrite crystals and coal fragments also occur in the mudstone bed.

Although it is difficult to compare the location of the first Maastrichtian vertebrate fossil site from the Rusca Montană Basin (CODREA et al., 2009, in press; FEIGI et al., 2010) to that of the site described here, the marked lithological differences between the fossiliferous deposits (red fine-grained overbank and gray channel sandstones for the site located “near Lunca Cernii de Sus”, respectively gray-blue overbank mudstones for the Fărcădeana Brook site, presented in this paper) suggest that the two fossil assemblages come different sedimentary bodies, representing distinct environments of the same fluvial system. Taking into account that the two sites also appear to sample different vertebrate groups (pending the discovery of further fossils), the differences in lithology and taxonomic composition are potential arguments supporting the idea that the fossil assemblage described herein comes from a different location (“site”) than the one described by Codrea and co-workers.




Figure 1. Location of the fossiliferous site from Fărcădeana Brook valley, Negoiu Village, Hunedoara County. Inset showing the location of the Rusca Montană area in Western Romania (Base map taken from Google Earth).

Figura 1. Amplasarea sitului fosilifer de pe valea Pârâului Fărcădeana, satul Negoiu, Județul Hunedoara. În medalion este ilustrată localizarea zonei Rusca Montană în partea de vest a României (Harta de bază preluată din Google Earth).

The anurans are the best represented group of microvertebrates recovered so far. Most of the specimens found are not diagnostic at a lower taxonomic level; these include a globular humeral ball (FGGUB v.548, Fig. 2b), typical anuran tibiofibular fragments (FGGUB v.551, Fig. 2a), and a fragmentary vertebra (FGGUB v.560). The tibiofibulae have the tibial and fibular components of similar size, wider at the epiphyses and narrowing to the diaphysis of the element. Distinct

sulci separate these components on both the anterior and posterior face of the tibiofibula, the former being more conspicuous. The only anuran remains determinable at specific level include a prearticular and an ilium, both incomplete. A small left ilium fragment (FGGUB v.550, Fig 2e) preserves the anterior part of the acetabular region, and the posterior part of the iliac crest. In lateral view, the acetabular region is fairly deep, bordered by a well-marked acetabular rim, prominent anteroventrally projecting beyond the ventral margin of the bone. Even if only a small posterior part of the iliac crest is preserved, this structure seems to have been rather tall. These features are not sufficient to positively assign v.550 to a certain taxon; nevertheless, they are reminiscent of those seen in ilia of *Paralatonia transylvanica* (VENCZEL & CSIKI, 2003), a common taxon in the Maastrichtian floodplain deposits of the Hațeg Basin, and also mentioned from the Transylvanian Basin at Oarda de Jos (JIPA et al., 2010). Three fragments of a large left prearticular (FGGUB v.549, Fig. 2c-d) have also been recovered. The largest of the three fragments preserves the area of the paracoronoid crest. The Meckelian groove is constricted at the level of the paracoronoid crest, being nevertheless well developed along all three fragments. The linguallly-projecting coronoid process has a concave dorsal surface. All these features are mentioned in the initial description of *Paralatonia*, supporting the referral of v.549 to this taxon.

The amphibians are also represented by a small dentary fragment, preserving the base of three thin cylindrical subpleurodont teeth, broken at the level of the dental parapet (FGGUB v.557, Fig. 2f). The Meckelian groove is closed; three oval foramina are present on the smooth labial side of the dentary. These features are typical for albanerpetontid lissamphibians (GARDNER, 2000), also present in the coeval deposits of the Hațeg (FOLIE & CODREA, 2005; GRIGORESCU et al., 1999) and the Transylvanian (CODREA et al., 2010a, 2010c; JIPA et al., 2010) basins, where they were referred to the genus *Albanerpeton* (ESTES & HOFFSTETTER, 1976).

Three crocodylian teeth have been recovered, belonging to two dental morphotypes. One of these (FGGUB v.555, Fig. 2g) has an equilateral triangle outline, is strongly labiolingually compressed and has a bulging, thicker central part, on the lingual face. One of the edges is broken, but the other bears a serrated carina, with large, subequal denticles, slightly decreasing in size from the tip to the base of the tooth, while concomitantly increasing in frequency. Two other teeth, also bearing serrated carinae, are subconical, taller than wide when compared to the first morphotype, apically recurved both distally and linguallly. The tips of both teeth are blunt, slightly worn; the wear facets are mesio-labially facing, longitudinally elongate ovals, reminiscent of those seen in teeth with comparable morphology from the Hațeg Basin. In FGGUB v.554 (Figs. 2h-i) the denticles decrease in size from the apex to the base. FGGUB v.556 (Figs. 2j-k) is smaller and has a lanceolate outline; it preserves part of the cylindrical root, and shows a slight constriction at the base of the crown. Similar dental morphotypes have been referred to the ziphodont crocodylian *Doratodon* (SEELEY, 1881), a terrestrial predator quite common in the Maastrichtian deposits of the Hațeg Basin (GRIGORESCU et al., 1999; MARTIN et al., 2006; VASILE & CSIKI, 2010) and in southwestern Transylvania (CODREA et al., 2010a, 2010b, 2010c; JIPA et al., 2010). The former morphotype was probably restricted to the distal part of the jaw, while the latter occupied a more anterior position.

Theropods are also represented by two isolated teeth with distinctive morphologies. One of these (FGGUB v.552, Figs. 2m-n) is subconical, long and narrow, labiolingually compressed and only slightly recurved mesodistally, bearing two unequal carinae. The mesial carena is smaller, migrated linguallly, and bears no denticles; it slowly diminishes towards the base of the crown where it ends in a small constriction. The distal carena is more conspicuous and bears clearly separated denticles, which decrease apically. The denticles are low and wide, with rounded tips, their axis being inclined toward the tooth apex. Faint longitudinal wrinkles are noticeable on both the lingual and the labial side. These features are similar to those described in the "*Richardoestesia*" morphotype (CURRIE et al., 1990), belonging to a small theropod dinosaur, a morphotype also reported from the microvertebrate assemblages of the Hațeg Basin (CODREA et al., 2002; VASILE & CSIKI, 2010). The second theropod tooth morphotype is represented by a recurved, labiolingually compressed tooth, markedly "D"-shaped in cross-section with asymmetrical lingual and labial sides (FGGUB v.553, Figs. 2o-p). The lingual side is flattened and shows a central concavity enclosed by the unserrated, linguallly projecting rim-like edges. The concavity is crossed by three longitudinal ridges, which divide it into four narrow and long depressions. The largest of the three ridges connects the tooth apex to the middle of its base. The remaining two ridges are shorter and fade towards the base and apex of the crown; the posterior ridge is more conspicuous than the anterior one. The convex labial side is also crossed by four longitudinal ridges separated by shallow depressions. Of these, the most prominent ridge is placed medially, equally distanced from the edges, with two smaller posterior, and one anterior ridge. This morphotype is similar to teeth assigned to *Paronychodon* (COPE, 1876) (e.g., SANKEY, 2008); only one single specimen of this taxon has been reported so far from Romania, from the Maastrichtian site of Totești-baraj (CODREA et al., 2002).

The lizards are represented by two fragmentary vertebrae, preserving only the centrum. These are procoelous and massive, ending in condyles slightly larger in diameter than the centrum itself. No haemal keel is present, the ventral side of the centrae being transversely convex and smooth. FGGUB v.559 (Fig. 2l) is longer and has a pair of foramina piercing the floor of the neural canal, placed near the midline of the centrum at about mid-length. FGGUB v.558 is thicker dorso-ventrally when compared to its length, and has four foramina on the ventral surface, asymmetrically placed in the right and left halves. The two foramina on the left side are small, and placed in a row lateral to the sagittal midline of the centrum. On the right side, one large foramen is placed near the midline, at about mid-length, while the second one, of approximately the same size, is migrated laterally and anteriorly. Similar foramina, placed in a similar irregular pattern, have been reported from the Hațeg Basin in vertebrae assigned to *Bicuspidon hatzegiensis* (FOLIE & CODREA, 2005); nevertheless, the presence of this feature is probably not sufficient to support the presence of the same taxon in Rusca Montană.




Figure 2. Microvertebrate fossil remains from Negoiu. a: FGGUB v.551 – Anuran tibiofibula fragment, anterior view; b: FGGUB v.548 – Anuran humeral proximal fragment; c-d: FGGUB v.549 – *Paralatonia transylvanica* left prearticular fragment, lingual and labial view; e: FGGUB v.550 – ?*Paralatonia transylvanica* left ilium fragment, lateral view; f: FGGUB v.557 – Albanerpetontid left dentary fragment, lingual view; g: FGGUB v.555 – *Doratodon* posterior tooth, lingual view; h-i: FGGUB v.556 – *Doratodon* anterior tooth, lingual and labial view; j-k: FGGUB v.554 – *Doratodon* anterior tooth, lingual and labial view; l: FGGUB v.559 – Partial lizard vertebra, ventral view; m-n: FGGUB v.552 – “*Richardoestesia*” morphotype, labial and lingual view; o-p: FGGUB v.553 – “*Paronychodon*” morphotype, lingual and labial view; q-r: FGGUB v.562 – Geckonoid eggshell fragment, inner and outer view; s-t: FGGUB v.563 – Indeterminate eggshell fragment, outer and inner view. Scale bars: 1 mm.

Figura 2. Resturi de microvertebrate fosile de la Negoiu. a: FGGUB v.551 – Fragment de tibiofibulă de anure, vedere anterioară; b: FGGUB v.548 – Fragment proximal humerus de anure; c-d: FGGUB v.549 – *Paralatonia transylvanica*, fragment de prearticular stâng, vedere linguală și labială; e: FGGUB v.550 – ?*Paralatonia transylvanica* fragment de ilion stâng, vedere laterală; f: FGGUB v.557 – Albanerpetontid, fragment de dentar stâng în vedere linguală; g: FGGUB v.555 – *Doratodon*, dinte posterior, vedere linguală; h-i: FGGUB v.556 – *Doratodon*, dinte anterior, vedere linguală și labială; j-k: FGGUB v.554 – *Doratodon*, dinte anterior, vedere linguală și labială; l: FGGUB v.559 – Vertebra parțială de șopârlă, vedere ventrală; m-n: FGGUB v.552 – Morfortipul “*Richardoestesia*”, vedere labial și linguală; o-p: FGGUB v.553 – Morfortipul “*Paronychodon*”, vedere linguală și labială; q-r: FGGUB v.562 – Fragment de coajă de ou de tip geckonoid, vedere internă și externă; s-t: FGGUB v.563 – Fragment nedeterminat de coajă de ou, vedere externă și internă. Scara grafică: 1 mm.

Two eggshell types have been recovered, represented by small, 2-5 mm long and wide fragments. The first morphotype, about 0.17 mm thick, bears a distinctive pattern of craters and small tubercles of various sizes on the outer surface (FGGUB v.562, Figs. 2q-r). The inner surface is smooth, the ends of the composing calcitic prisms being tightly packed. The dispersituberculate pattern from the outer surface of these eggshells is consistent with that seen in the geckonoid eggshell morphotype (MIKHAILOV, 1991). Eggshells of this type have been reported from Maastrichtian deposits of both the Hațeg (e.g. CODREA et al., 2002; CSIKI et al., 2008) and Transylvanian (CODREA et al., 2010a, b, c; JIPA et al., 2010) basins. The second morphotype is represented by slightly thicker (0.37-0.44 mm) eggshells with a prismatic structure producing a smooth inner surface and an outer surface uniformly covered by small, sub-equal, tightly packed tubercles (FGGUB v.563, Figs. 2s-t). A more detailed analysis is needed to thoroughly assess the affinities of these eggshells; thin eggshell fragments different from the geckonoid morphotype have also been mentioned from the Hațeg Basin (e.g. CODREA et al., 2002; CSIKI et al., 2008), but their presumed theropod origin is yet to be supported.

The most abundant fossils from the site are the gastropods, most of these similar to morphotypes reported previously from the Hațeg Basin by PANĂ et al. (2001); although several of the gastropods were mentioned as new taxa in that preliminary paper, no formal description of them is available, and thus they will be referred herein to as morphotypes, and not nominal taxa.

The largest part of the gastropod fossils from Negoiu consists of circular opercula (Figs. 3a-b), identical to the ones commonly encountered in many sites of the Hațeg Basin, and considered to belong to cyclophorids (PANĂ et al., 2001). The terrestrial cyclophorids are also represented by *Ischurosoma filholi* (Fig. 3c) and *Ajkaia* cf. *gregaria* (Fig. 3d); these are associated with fragmentary remains that suggest the presence of helicids (*Helicella*; Fig. 3e). On the other hand, the aquatic taxa appear to be represented by the lymneids *Lymnaea* (Fig. 3f) and *Physa* (Fig. 3g), as well as the planorbids *Planorbis planorbis* (Fig. 3h) and the “*Palaeoanisis septemgiratus*” morphotype (Figs. 3i-j).

The co-occurrence of terrestrial and aquatic gastropods within the same assemblage was already noted in several sites of the Hațeg Basin (PANĂ et al., 2001), and was used to suggest a mixed assemblage with parautochthonous terrestrial taxa (cyclophorids, helicids) washed into the depositional area from the surrounding land, along with largely autochthonous aquatic taxa characteristic for small-sized stagnant water bodies (planorbids, lymneids). The abundance of gastropod opercula, representing the most resistant part of the shell, is consistent with this idea, as the robust opercula might have been transported into the depressionary setting much easily, i.e. in larger numbers and better preservation state, than the more fragile shells.

A fairly large number of minute angiosperm fructifications have been extracted from the sample yielding the microvertebrate and gastropod remains. Most of them show morphologies similar to fruits reported from the Budurone microvertebrate fossil site, in the Hațeg Basin (MAY LINDFORS et al., 2010), considered to represent new taxa, but not yet described formally. One of the morphological types recovered from Negoiu (Fig. 4a) is similar to Taxon A of MAY LINDFORS et al. (2010: Fig. 2a), an ellipsoidal fruit probably related to the Normapolles group, similar to the Normapolles genus *Antiquocarya*. Other specimens are either capsular (Figs. 4b-d) or drupaceous (Figs. 4e-g) indehiscent fruits. Two different morphotypes of capsular, five-locular fruits are also similar to the ones mentioned from Budurone. One of them is ovoid (Fig. 4b) and the other circular (Figs. 4c-d); these have been referred to the Ericales (Taxon 10 and Taxon 11 of MAY LINDFORS et al., 2010), a common taxon in the European Upper Cretaceous. The drupaceous fruits are unilocular, one of the morphs found in Negoiu being reminiscent of those from Budurone (Fig. 4e). A further morphotype, represented by simple ovoidal structures (Fig. 4h), is also recorded in the Budurone fossil assemblage, but its affinities remain unclear. The present carpoflora confirms the presence of angiosperms belonging to the Normapolles group, documented previously by palynological studies from the Rusca Montană Basin (ANTONESCU et al., 1983), and augments the previously known floral list based mainly on leaf impressions (e.g. GIVULESCU 1966, 1968; PETRESCU & DUȘA, 1970, 1980, 1985; TUZSON, 1913). This mesoflora assemblage is similar in size and abundance of the taxa to other Late Cretaceous floras from Europe. Endozoochory has been postulated as an important dispersal mechanism in Late Cretaceous angiosperms with drupaceous fruits (ERIKSSON et al., 2010), a possible vector for such dispersal in the Rusca Montană Basin being the multituberculate mammals, already reported by CODREA et al. (2009, in press).

CONCLUSIONS

This paper represents the first record of several non-mammalian microvertebrates - anurans, albanerpetontids, lizards, the ziphosuchian *Doratodon*, the theropods “*Paronychodon*” and “*Richardoestesia*” -, along with fragmentary eggshells, gastropods, and plant mesofossils, from the Maastrichtian deposits of the Rusca Montană Basin. These finds add to the previously reported fossil taxa (CODREA et al., 2009, in press; FEIGI et al., 2010; PETRESCU & DUȘA, 1985), thus expanding the previously known taxon list (Table 1) and contribute to the reconstruction of the biodiversity of the Romanian Maastrichtian ecosystems, while also extending the known palaeogeographical distribution range of these taxa. The presence of similar microvertebrates, gastropods and plant mesofossils in the Maastrichtian assemblages from the Hațeg and Rusca Montană basins lends further support for a synchronous sedimentation, and coeval ecosystems, in the two neighbouring areas, already suggested by lithological, palynological and paleofloristic data. More than that, the identification of similar assemblages in different sites spread over the Hațeg, Rusca Montană and Transylvanian basins could suggest a biostratigraphical importance for some of the microvertebrates, gastropods and mesofloristic taxa in the Romanian Maastrichtian.




Figure 3. Fossil gastropods from Negoiu (uncatalogued specimens). Fam. Cyclophoridae: a, b - Cyclophorid operculae; c - *Ischurosoma filholi*; d - *Ajkaia cf. gregaria*; Fam. Helicidae: e - *Helicella* sp.; Fam. Lymnaeidae: f - *Lymnaea* sp.; g - *Physa* sp.; Fam. Planorbidae: h - *Planorbis planorbis*; i, j - "*Palaeoanisis septemgiratus*". Scale bar: 1 mm.

Figura 3. Gastropode fosile de la Negoiu (specimene neînregistrate). Fam. Cyclophoridae: a, b - Operculi de cyclophoride; c - *Ischurosoma filholi*; d - *Ajkaia cf. gregaria*; Fam. Helicidae: e - *Helicella* sp.; Fam. Lymnaeidae: f - *Lymnaea* sp.; g - *Physa* sp.; Fam. Planorbidae: h - *Planorbis planorbis*; i, j - "*Palaeoanisis septemgiratus*". Scara grafică: 1 mm.




Figure 4. Fossil angiosperm fructifications from Negoiu (uncatalogued specimens). a: Indehiscent fruit related to the Normapolles group; b-d: Five-locular capsules related to Ericales; e-g: Indeterminate drupaceous fruits; h: Ovoidal morphotype of unclear affinities. Scale bar: 1 mm.

Figura 4. Fructificații fosile de angiosperme de la Negoiu (specimene neînregistrate). a: Fruct indehiscent apropiat de grupul Normapolles; b-d: Capsule pentaloculare apropiate de Ericales; e-g: Fructe drupacee nedeterminate; h: Morfotip ovoidal cu afinități neclare. Scara grafică: 1 mm.

Table 1. List of the Maastrichtian vertebrate taxa (including eggshells) from the Rusca Montană Basin, based on finds from Negoiu, Fărcădeana Brook (this paper) and near Lunca Cernii de Sus (CODREA et al., 2009, in press; FEIGI et al., 2010).

Tabel 1. Lista vertebratelor (inclusiv coji de ouă) maastrichtiene din Bazinul Rusca Montană, pe baza descoperirilor de la Negoiu, Pârâul Fărcădeana (lucrarea de față) și din apropiere de Lunca Cernii de Sus (CODREA et al., 2009, in press; FEIGI et al., 2010).

Taxa/Morphotypes	Location	Element
AMFIBIA		
Allocaudata		
Albanerpetontidae ? <i>Albanerpeton</i>	Negoiu, Fărcădeana Brook	dentary fragment
Anura		
Discoglossidae <i>Paralatonia transylvanica</i>	Negoiu, Fărcădeana Brook	ilium, prearticulary
REPTILIA		
Chelonia		
Cryptodira ? <i>Kallokibotion bajazidi</i>	near Lunca Cernii de Sus	carapace fragments
Lepidosauria		
Sauria indet.	Negoiu, Fărcădeana Brook	2 vertebrae
Archosauria		
?Eusuchia (? <i>Allodaposuchus</i>)	near Lunca Cernii de Sus	ilium, ischium, pubis, 2 caudal vertebrae
Ziphosuchia <i>Doratodon</i>	Negoiu, Fărcădeana Brook	3 isolated teeth
Dinosauria		
Theropoda		
Theropoda indet.	near Lunca Cernii de Sus	isolated tooth
Velociraptorinae indet.	near Lunca Cernii de Sus	3 isolated teeth
„ <i>Richardoestesia</i> ”	Negoiu, Fărcădeana Brook	tooth
„ <i>Paronychodon</i> ”	Negoiu, Fărcădeana Brook	tooth
Sauropoda		
Titanosauridae indet.	near Lunca Cernii de Sus	caudal vertebra
Ornithopoda		
<i>Zalmoxes</i>	near Lunca Cernii de Sus	2 teeth, caudal vertebra
MAMMALIA		
Multituberculata		
Cimolodonta Kogaionidae indet.	near Lunca Cernii de Sus	2 teeth (P2, p4)
VETROVATA		
Geckonoid morphotype	Negoiu, Fărcădeana Brook	eggshell fragments
Indeterminate morphotype	Negoiu, Fărcădeana Brook	eggshell fragments

The sediment yielding the microvertebrates, gastropods and fruits described herein is gray-blue, a colour indicative for a poorly oxygenated depositional environment, as is the presence of coal fragments and pyrite crystals (e.g., WHITE et al., 1990; HAMILTON-TAYLOR & DAVISON, 1995). Together with the fine-grained sediments, they suggest that sedimentation took place in a poorly-drained floodplain, with ponds and marshy areas (e.g., WRIGHT et al., 2000). The preservation of fragile fossil structures, such as the thin calcitic eggshell fragments and gastropod shells, alongside buoyant and therefore easily transportable structures like small-sized fruits, suggests that they did not undergo a significant amount of transport, another argument supporting the hypothesis of a hydrodynamically quiet, poorly drained floodplain environment.

Many features of the depositional environment and fossil record from the newly reported Negoiu site are similar to those described in the Budurone microvertebrate fossil site from the Hațeg Basin (CSIKI et al., 2008), and lend further support to the identification of the depositional setting as a poorly drained, water-logged area, inhabited by the semi-aquatic invertivore vertebrates and freshwater gastropods, the top of the local food chain being represented by small theropods like *Richardoestesia*, or by ambush-preying terrestrial crocodiles (*Doratodon*), feeding on the semi-aquatic (anurans, albanerpetontid) or terrestrial (lizards) taxa that either lived in this area or visited it for feeding or drinking. The additional terrestrial gastropod remains and the eggshell fragments were washed into the site from the surrounding emergent areas, while the angiosperm fruits suggest the presence of a thick canopy surrounding the water body.

ACKNOWLEDGEMENTS

The fieldwork and part of the laboratory preparation was supported by, and is a contribution to, the CNCSIS Grant 1930/2009. ȘV also benefited from POSDRU/88/1.5/S/61150 "Doctoral studies in the field of Life and earth sciences", project co-financed through Sectorial Operational Program for the Development of Human Resources 2007-2013 from European Social Fund. Fellow geologist Bogdan Săvescu, as well as Mr. Dumitru Huzoni and the family of Mr. Moise Brăilă are warmly thanked for contributing with technical and logistic support.

The reviewers, Pascal Godefroit (Royal Institute of Natural Sciences, Bruxelles) and Vlad Codrea (University Babeș-Bolyai, Cluj-Napoca), are thanked for their constructive comments and suggestions that helped improve the initial version of the text. Vlad Codrea is also thanked for pointing out, and sending the authors, additional literature.

REFERENCES

- ANTONESCU E., LUPU DENISA, LUPU M. 1983. *Corrélacion palinologică du Crétacé terminal du sud-est des Monts Metaliféri et des Dépressions de Hațeg et de Rusca Montană*. Anuarul Institutului de Geologie și Geofizică. București **59**: 71-77.
- BITOIANU CORNELIA. 1970. *Observații asupra constituției petrografice a cărbunilor de la Rusca Montană (Județ Caraș-Severin)*. Institutul Geologic. Studii tehnice și economice. Seria A : Prospectiuni și explorări geologice. București. **8**: 105-120.
- CANTUNIARI ȘT. 1937. *Études géologiques dans les Monts Poiana Ruscă. I. Bassin de Rusca. Région de Rusca Montană*. Comptes Rendus des Séances de l'Institut Géologique de Roumanie. București. **21**: 156-168.
- COPE E. D. 1876. *Descriptions of some vertebrate remains from the Fort Union beds of Montana*. Proceedings of the Academy of Natural Sciences of Philadelphia: 248-261.
- CODREA V., SMITH T., DICA P., FOLIE ANNELISE, GARCIA GÉRALDINE, GODEFROIT P., VAN ITTERBEECK J. 2002. *Dinosaur egg nests, mammals and other vertebrates from a new Maastrichtian site of Hațeg Basin (Romania)*. Comptes Rendus Palevol. Elsevier. Amsterdam. **1**: 173-180.
- CODREA V., GODEFROIT P., SMITH T., JIPA-MURZEA C. 2009. *Maastrichtian land vertebrates in Rusca Montană Basin (Romania)*. In: Godefroit & Lambert (Eds.) Tribute to Charles Darwin and Bernissart iguanodonts: New Perspectives on Vertebrate Evolution and Early Cretaceous Ecosystems: 29.
- CODREA V., BARBU O., JIPA-MURZEA C. 2010a. *Upper Cretaceous (Maastrichtian) landvertebrate diversity in Alba District (Romania)*. Bulletin of the Geological Society of Greece, Proceedings of the 12th International Congress. **63**(2): 594-601.
- CODREA, V., JIPA-MURZEA C., CSIKI Z., BARBU O. 2010b. *Maastrichtian dinosaurs in SW Transylvania (Romania)*. Scientific Annals, School of Geology, Aristotle University of Thessaloniki. Proceedings of the XIX CBGA Congress. Special volume. **99**: 69-74.
- CODREA V., VREMIR M., JIPA C., GODEFROIT P., CSIKI Z., SMITH T., FĂRCAȘ CRISTINA. 2010c. *More than just Nopcsa's Transylvanian dinosaurs: A look outside the Hațeg Basin*. Palaeogeography, Palaeoclimatology, Palaeoecology. Elsevier. Amsterdam. **293**: 391-405.
- CODREA V., GODEFROIT P., JIPA C., SMITH T. (in press). *First find of Maastrichtian terrestrial vertebrates in Rusca Montană Basin*. In: Godefroit (Ed.) Bernissart. Dinosaurs in Depth. Indiana University Press, Bloomington.
- CSIKI Z., IONESCU ANA, GRIGORESCU D. 2008. *The Budurone microvertebrate site from the Maastrichtian of the Hațeg Basin – flora, fauna, taphonomy and paleoenvironment*. Acta Palaeontologica Romaniaae. Edit. Universității „Al. I. Cuza”. Iași. **6**: 49-66.
- CURRIE P. J., RIGBY JR. J. K., SLOAN R. E. 1990. *Theropod teeth from the Judith River Formation of Southern Alberta, Canada*. In: Carpenter & Currie (Eds.) Dinosaur Systematics: Perspective and Approaches. Cambridge University Press. Cambridge: 107-125.
- DINCĂ A. 1977. *Geologia Bazinului Rusca Montană. Partea de vest*. Anuarul Institutului de Geologie și Geofizică. București. **52**: 99-173.
- DINCĂ A., TOCORJESCU MARIA, STILLA A. 1972. *Despre vârsta depozitelor continentale cu dinozaurieni din Bazinele Hațeg și Rusca Montană*. Dări de Seamă ale Institutului Geologic. București. **58**(4): 83-94.
- DUȘA A. 1970. *Contribuții la studiul petrografic al cărbunilor din Bazinul Rusca Montană*. Studia Universitatis “Babeș-Bolyai”. Series Geologia-Mineralogia. Cluj-Napoca. **15**(1): 33-41.
- DUȘA A. 1974. *Aspecte ale formării cărbunilor din Bazinul Rusca Montană*. Studia Universitatis “Babeș-Bolyai”. Series Geologia-Mineralogia. Cluj-Napoca. **2**: 36-42.
- DUȘA A. 1987. *Zăcămintele de huile din Cretacicul superior*. In: Petrescu, Nicorici, Bițoianu, Țicleanu, Todros, Ionescu, Mărgărit, Nicorici, Dușa, Patruțoiu, Munteanu & Buda (Eds.) Geologia zăcămintelor de cărbuni. **2**: 74-81. Editura Tehnică, București.
- DUȘA A. & BĂRILĂ M. 1973. *Aspecte petrografice și paleobotanice ale cărbunilor din Bazinul Rusca Montană*. Studia Universitatis “Babeș-Bolyai”. Series Geologia-Mineralogia. Cluj-Napoca. **18**(1): 31-38.
- ERIKSSON O., FRIIS ELSE MARIE, LÖFGREN P. 2000. *Seed size, fruit size and dispersal spectra in angiosperms from the Early Cretaceous to the Late Tertiary*. American Naturalist. University of Chicago Press. Chicago, Illinois. **156**(1): 47-58.
- ESTES R. & HOFFSTETTER R. 1976. *Les Urodèles du Miocène de La Grive-Saint-Alban (Isère, France)*. Bulletin du Muséum National d'Histoire Naturelle, 3^e série, no. 398, Sciences de la Terre. Paris. **57**: 297-343.
- FEIGI Ș. V., JIPA C., SOLOMON A. 2010. *Paleomedii maastrichtiene în Bazinul Rusca Montană*. Lucrările celui de-al X-lea Simpozion Național Studentesc „Geoecologia”, Petroșani. **1**: 33-36.
- FOLIE ANNELISE & CODREA V. 2005. *New lissamphibians and squamates from the Maastrichtian of Hațeg Basin, Romania*. Acta Palaeontologica Polonica. Warsaw. **50**(1): 57-71.

- GARDNER J. D. 2000. *Revised taxonomy of albanerpetontid amphibians*. Acta Palaeontologica Polonica. Warsaw. **28**(1-2): 103-121.
- GIVULESCU R. 1966. *Sur quelques plantes fossiles du Danien de Roumanie*. Comptes Rendus de l'Académie des Sciences de Paris. **262**(D): 1933-1936.
- GIVULESCU R. 1968. *Nouvelles plantes fossiles du Danien de Roumanie*. Comptes Rendus de l'Académie des Sciences de Paris. **267**(D): 880-882.
- GRIGORESCU D., HARTENBERGER J.-L., RĂDULESCU C., SAMSON P., SUDRE J. 1985. *Découverte de mammifères et dinosaures dans le Crétacé supérieur de Pui (Roumanie)*. Comptes Rendus de l'Académie des Sciences de Paris. **301**: 1365-1368.
- GRIGORESCU D. 1992. *Nonmarine Cretaceous Formations of Romania*. In: Matter & Chen (Eds.) Aspects of Nonmarine Cretaceous Geology. China Ocean Press. Beijing: 142-164.
- GRIGORESCU D. 2010. *The Latest Cretaceous fauna with dinosaurs and mammals from the Hațeg Basin – A historical overview*. Palaeogeography, Palaeoclimatology, Palaeoecology. Elsevier. Amsterdam. **293**: 271-282.
- GRIGORESCU D., VENCZEL M., CSIKI Z., LIMBEREA R. 1999. *New microvertebrate fossil assemblages from the Uppermost Cretaceous of the Hațeg Basin (Romania)*. Geologie en Mijnbouw. Utrecht. **78**: 301-314.
- HAMILTON-TAYLOR J. & DAVISON W. 1995. Redox-driven cycling of trace elements in lakes. In Lerman, Imboden & Gat (Eds.) Physics and chemistry of lakes (second edition). Springer-Verlag. New York: 217-264.
- JIPA C., SOLOMON A., FEIGI Ș. V., MICLEA ANGELA. 2010. *Microvertebrate continentale maastrichtiene de la Oarda de Jos (jud. Alba) – Date preliminare*. Lucrările celui de-al X-lea Simpozion Național Studentesc „Geoecologia”. Petroșani. **1**: 41-44.
- MAY LINDFORS SANDRA, CSIKI Z., GRIGORESCU D., FRIIS ELSE MARIE. 2010. *Preliminary account of plant mesofossils from the Maastrichtian Budurone microvertebrate site of the Hațeg Basin, Romania*. Palaeogeography, Palaeoclimatology, Palaeoecology. Elsevier. Amsterdam. **293**: 353-359.
- MAMULEA A. 1955. *Cercetări geologice în regiunea Rusca Montană – Lunca Cernii (comunicare preliminară)*. Dări de Seamă ale Ședințelor Comitetului Geologic. București. **39**: 172-178.
- MARTIN J. E., CSIKI Z., GRIGORESCU D., BUFFETAUT E. 2006. *Late Cretaceous crocodylian diversity in Hațeg Basin, Romania*. Hantkeniana. Eötvös Loránd Tudományegyetem. Budapest. **5**: 31-37.
- MIKHAILOV K. E. 1991. *Classification of fossil eggshells of amniotic vertebrates*. Acta Palaeontologica Polonica. Warsaw. **36**(2): 193-238.
- NOPCSA F. 1905. *Zur Geologie der Gegend zwischen Gyulafehérvár, Déva, Ruszkabánya und der Rumänischen Landesgrenze*. Mitteilungen aus dem Jahrbuche der Königlich Ungarischen Geologischen Reichsanstalt, Budapest. **14**: 93-279.
- PANĂ IOANA, GRIGORESCU D., CSIKI Z., COSTEA C. 2002. *Paleo-ecological significance of the continental gastropod assemblages from the Maastrichtian dinosaur beds of the Hațeg Basin*. Acta Palaeontologica Romaniae. Editura Vasiliana 98. Iași. **3**: 337-343.
- PETRESCU I. & DUȘA A. 1980. *Asupra unui nou punct paleofloristic din Cretacicul superior al Bazinului Rusca Montană*. Buletinul Societății de Științe Geologice din R. S. România. București. **12**: 165-172
- PETRESCU I. & DUȘA A. 1980. *Flora din Cretacicul superior de la Rusca Montană – o raritate în patrimoniul paleobotanic național*. Ocrotirea naturii și a mediului înconjurător. București. **24**(2): 147-155.
- PETRESCU I. & DUȘA A. 1985. *Paleoflora din Senonianul Bazinului Rusca Montană*. Dări de Seamă ale Institutului de Geologie și Geofizică. București. **69**(3): 107-124.
- POP GH. & PETRESCU I. 1983. *Considerații paleoclimatice asupra vegetației din Cretacicul superior de la Rusca Montană*. Studia Universitatis “Babeș-Bolyai”. Series Geologia-Geographia. Cluj-Napoca. **28**: 49-54.
- SANKEY JULIA T. 2008. *Diversity of Latest Cretaceous (Late Maastrichtian) Small Theropods and Birds: Teeth from the Lance and Hell Creek Formations, USA*. In: Sankey & Baszio (Eds.) Vertebrate Microfossil Assemblages: their role in paleoecology and paleobiogeography. Indiana University Press. Bloomington: 117-134.
- SCHAFARZIK F. 1906. *A krassószörényi Pojána-Ruszkahegység DNY-i részének geológiai viszonyai*. A Magyar Királyi Földtani Intézet évi Jelentése 1905-ről. Franklin-Társulat Könyvnyomdája. Budapest. 84-95.
- SEELEY H. G. 1881. *The reptile fauna of the Gosau Formation preserved in the Geological Museum of the University of Vienna*. Quarterly Journal of the Geological Society of London. **37**: 620-704.
- STRUTINSKI C. & HANN H. P. 1986. *Reconsidération de la structure géologique de Rusca Montană et ses implications sur la tectonique du Massif de Poiana Ruscă*. Dări de Seamă ale Ședințelor Institutului de Geologie și Geofizică. București. **70-71**(5): 255-268.
- TUZSON J. 1913. *Adatok Magyarország fosszilis flórájához (Addindamenta ad floram fossilem Hungariae III.)*. A Magyar Királyi Földtani Intézet Évkönyve. Franklin-Társulat Könyvnyomdája. Budapest. **21**(8): 209-233.
- VASILE Ș. & CSIKI Z. 2010. *Comparative paleoecological analysis of some microvertebrate fossil assemblages from the Hațeg Basin, Romania*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **26**(1): 315-322.
- VENCZEL M. & CSIKI Z. 2003. *New frogs from the latest Cretaceous of Hațeg Basin, Romania*. Acta Palaeontologica Polonica. Warsaw. **48**(4): 609-616.

- WHITE T. S., MORRISON J. L., KUMP L. R. 1990. *Formation of iron sulfides in modern salt marsh sediments (Wallops Island, Virginia)*. In Orr & White (Eds.) *Geochemistry of sulfur in fossil fuels*. American Chemical Society Symposium Series. Oxford University Press. Oxford. **429**: 204-217.
- WRIGHT V. P., TAYLOR K. G., BECK V. H. 2000. *The paleohydrology of Lower Cretaceous seasonal wetlands, Isle of Wight, Southern England*. *Journal of Sedimentary Research*. Boulder, Colorado. **70**(3): 619-632.

Vasile Ștefan

University of Bucharest, Faculty of Geology and Geophysics, Laboratory of Paleontology,
1 N. Bălcescu Ave., RO-010041, Bucharest, Romania
E-mail address: yokozuna_uz@yahoo.com

Csiki Zoltán

University of Bucharest, Faculty of Geology and Geophysics, Laboratory of Paleontology,
1 N. Bălcescu Ave., RO-010041, Bucharest, Romania
E-mail address: zoltan.csiki@g.unibuc.ro

Received: March 30, 2011

Accepted: July 15, 2011