

## NEW CONTRIBUTIONS TO THE CIOCADIA MIDDLE MIOCENE FLORA (PART TWO)

PARASCHIV Valentin

**Abstract.** Further revisions and detailed determinations are made for the plant fossil assemblages discovered in the Middle Miocene deposits of Ciocadia village, South Carpathian Foredeep. Arcto-Tertiary geofloristic elements are certified, namely *Berberis* sp., *B. coloradensis*, *Ulmus pyramidalis*, *U. pardschlugiana*, and *Zelkova zelkovifolia*. The macroremains of these species are thoroughly described, figured and comparative analysis are made taking into account all the analogous fossils reported in Romania. Taphonomic and palaeoecological considerations are assumed to clarify the sedimentation conditions and factors and the palaeoenvironmental significance of the fossil plant occurrence is given.

**Keywords:** taxonomy, plant megafossils, *Berberis coloradensis*, Ciocadia, Middle Miocene.

**Rezumat. Noi contribuții la flora miocen medie de la Ciocadia (partea a doua).** Elemente din asociația de plante fosile descoperite în depozitele Miocenului mediu din aria localității Ciocadia (Avanfosa Carpaților Meridionali) sunt descrise detaliat și au fost supuse unor revizii. Provincia geofloristică Arctoterțiară (care domină astăzi continentul Nord-American, Asia de Est extratropicală și Europa) este reprezentată în flora fosilă de la Ciocadia prin *Berberis* sp., *B. coloradensis*, *Ulmus pyramidalis*, *U. pardschlugiana* și *Zelkova zelkovifolia*. Fosilele macroscopice de plante sunt analizate paleontologic, fotografiate și discutate comparativ în relație cu celelalte descoperiri similare din România. De asemenea, au fost simulate condițiile tafonomice și paleoecologice în care s-a realizat conservarea materialului vegetal sub formă de fosile și sunt oferite informații despre mediul depozițional (de sedimentare) și configurația paleoreliefului din timpul miocenului.

**Cuvinte cheie:** taxonomie, plante fosile, *Berberis coloradensis*, Ciocadia, Miocenul mediu.

### INTRODUCTION

Occurrences of Berberidaceae are rare in the area of the Paratethys Basin; instead Ulmaceae remains are recovered in numerous sites, as samaras and sometimes as foliage. Now we undertake detailed analysis of two peculiar specimens of *Berberis*, two distinct groups of fruits belonging to *Ulmus*, and there also is described, for the first time in the palaeoflora of Ciocadia, *Zelkova* foliage. Recent studies of some selected taxa of Ciocadia flora contributed to the revision (PARASCHIV, 2008a, b; PARASCHIV & SEBE, 2011) of the palaeofloristic assemblage. We focus on the rectification of the previous taxonomic data in order to attempt a more precise floristic correlation with other Romanian and European Miocene sites. The palaeofloristic assemblage of Ciocadia is of great stratigraphical importance because of independent dating of the fossiliferous deposit by small marine invertebrate remains belonging to the group of pteropods (particularly the shelled pteropod *Limacina*, see Pl. I, Fig. 11). These deposits correspond to the 'Spirialis Marls' (POPESCU, 1953; TUDOR, 1955; MARINESCU, 1969; HUICĂ, 1994; PARASCHIV, 2006) which consist in a bloom (associated with a red tide phenomenon) of fossil holoplanktonic gastropods with thin aragonitic shells, very fragile, and which is easily subject to dissolution. The depth of the water column may be approximated as a reflexion of the presence of upwelling currents which bring nutrient-rich bottom waters up into the photic zone. Such high level of nutrients would allow primary productivity to occur at an explosive rate (SCHOPF, 1980). Meanwhile, only an upwelling during Ciocadia's marls deposition, and a possible red tide could explain the mass occurrence of *Limacina* (= *Spiratella* (*Spirialis*) horizon). *Spiratella* horizon is characteristic to the Late Badenian (Kossovian) normal marine deposits in the Central Paratethys (IORGULESCU, 1952; BOHN-HAVAS et al., 2004). Pteropod taphocoenosis can be, also, generated by deposition of storm currents (VELCESCU, 1997). But the recognized large scale extension of *Spiratella* horizon may exclude a tempestitic hypothesis. These deposits are conformably overlain by marine, brackish water laminated grey marls (also rich in phytoclast assemblages), silty clays, sand and gravel beds belonging to the Early Sarmatian (Volhynian) stage. This complete series of shallow water deposits (Late Badenian-Early Sarmatian) provides a local to regional record of the flora, mollusc, insect and fish fauna (PARASCHIV, 2006).

### MATERIAL AND METHODS

For analytical observations and studies of the fossil remains a Carl Zeiss - Jena Technoval Stereo Microscope (Binocular) was used, followed by examination of the literature. The repository place for the studied specimens is the National Museum of Geology of the Geological Institute of Romania, Bucharest. The accepted morphological terminology used is mainly based on HICKEY (1979) and WING et al. (1999). Systematic organization and taxonomic terminology in this article are based on the works of KUBITZKI (1993) and TAKHTAJAN (2009).

## SYSTEMATIC PALAEOBOTANY

Phylum *Magnoliophyta* CRONQUIST 1996  
 Class *Magnoliopsida (Dicotyledons)* BRONGNIART 1843  
 Order *Berberidales* DUMORTIER 1829  
 Family *Berberidaceae* JUSSIEU 1789  
 Genus *Berberis* LINNAEUS 1753  
*Berberis* sp.  
 Pl. I, Figs. 1, 2.

Material: BCI.0322, BCI.0322a.

Description. Incomplete obovate leaf, coriaceous, slightly asymmetrical, with missing base and rounded apex; more than 35 mm in length and 14 mm is the maximum width. The lower part of the lamina is long, attenuated to cuneate. Leaf margin serrate, with tiny spines teeth, or irregularly with crenulated aspect, conspicuously revolute. Midvein strong, raised adaxially, slightly curved; secondary veins of festooned brochidodromous type, conspicuous, diverging at an angle of 20° to 35° from base to apex, and looping with upper secondary veins; inter-secondary veins simple, straight or arched, fusing with secondary veins; tertiary veins reticulate and quaternary veins forming polygonal areoles. The leaf was severely damaged during diagenesis being transversely ripped apart in close range; nevertheless the still preserved leaf architecture maintains well defined characters that relate it to the genus *Berberis*. The most probable resemblance of our specimen is with fossil *Berberis berberidifolia* (HEER) PALAMAREV & PETKOVA (PALAMAREV & PETKOVA, 1987; KVAČEK et al., 2004).

Remarks. *Berberis* remains are cited from the Palaeocene to Pleistocene sediments of the Northern Hemisphere (PETRESCU & DRAGASTAN, 1981; LI et al., 2010). In Romania, it is described *Berberis* sp. from the **Cornești-Aghireș** (Cluj County, PETRESCU et al., 1997) Oligocene (Late Rupelian-Early Chattian) deposits, *Berberis* sp. from the **Coruș** (Cluj County, ȚICLEANU & GIVULESCU, 1978) Early Miocene deposits, *B. andreanszkyi* KVAČEK & ERDEI from the Early Sarmatian deposits of the **Morilor Valley** section (Mehedinți County, PARASCHIV 2006, 2008b), *Berberis* sp. from **Corni** (Neamț County, ȚICLEANU & MICU, 1979) in the Middle Miocene (Middle Sarmatian-Basarabian) deposits, *Berberis kymeana* (UNGER) KVAČEK & ERDEI from **Valea Neagră de Criș** (Valea Crișului, ante Valea Neagră, Bihor County) in the Early Pannonian deposits (GIVULESCU, 2002), *Berberis goinai* GIVULESCU, *Berberis inopinata* GIVULESCU, *Berberis* cf. *mougeoti* (HEER) GIVULESCU and *Berberis lanceolata* GIVULESCU are described from the **Chiuzbaia** palaeoflora (Maramureș County) in the Late Miocene deposits (Late Pontian = Pannonian s.l. G/H, GIVULESCU, 1990). *Berberis* of Cornești-Aghireș flora show a pinnately brochidodromous venation, and leaf margin irregularly toothed with tiny oblique teeth. There is no resemblance with any of Ciocadia specimens and it may be close to *B. lycium* Royle (see LI et al., 2010) of the section Asiaticae. ȚICLEANU & GIVULESCU (1978) believed to recognize another specimen of *Berberis* in Coruș beds. They only give few characters like ‘oblong leaf impression (40 mm long, 9 mm wide) with cuneate base and a 2 mm long petiole, rounded apex and entire margin; venation distinct’, unfortunately without a helpful illustration. The presence of *Berberis* in the Sarmatian sediments of Corni is suitable but the leaf fragment lack in details. This poorly preserved specimen shows a weak brochidodromous venation that can fit with a question mark to the Berberidaceae. Although the fossil record is relatively abundant (especially in the Central and North American Tertiary floras), LI et al. (2010) validate only seventeen true species of fossil *Berberis* preserved as leaves and omitted two other species described by GIVULESCU (1990) in Chiuzbaia palaeoflora, respectively *Berberis inopinata* and *Berberis lanceolata*. The extant *Berberis vulgaris* Linnaeus is the single species of *Berberis* which grow naturally in Romania, as shrubs and bushes in sunny and dry hills, on rocky substrates (SĂVULESCU et al., 1955; CIOCĂRLAN, 2009). It is native to continental Europe, except for the northwest, and is rare in the Mediterranean region.

*Berberis coloradensis* AXELROD 1987

Pl. I, Fig. 3.

*Mahonia hakeaeformis* BECKER; BECKER, 1969, p. 90, Pl. XXVI, Fig. 4.

*Berberis coloradensis* AXELROD; AXELROD, 1987, p. 110, Pl. XXII, Figs. 1, 2, 3, 6, 7.

Material: BCI.0181.

Description. Small leaf almost complete, simple, narrowly elliptic, with the base quasi-symmetric, cuneate, basal angle acute, apex absent; the length is of about 20 mm and width of 6 mm; midrib firm, thin in the most part of the lamina and thickened in the base, where it becomes slightly swollen and divergent (near the node around which the leaf can flex); petiole absent; basal acrodromous venation with two lateral secondary veins rising at low angles forming elongated and asymmetric loops, of characteristic festooned brochidodromous type; other secondaries originate at angles of 70°-80° extending straight to form brochidodromous loops that are elongated parallel to midrib joining to it; tertiary veins seems to be randomly reticulate; margin sparsely dentate, with one prominent, spiny tooth near the middle of blade and tiny apiculate teeth irregularly placed.

Remarks. This form is represented by one poorly preserved specimen; however the distinctive acrodromous venation, the festooned brochidodromy of the secondaries, rare spinose teeth or the missing petiole justified the type suite of the designated species. *Berberis coloradensis* was compared by AXELROD (1987) to the extant *B. candidula* SCHNEIDER which is endemic to China (where it lives at the altitude of 1200-3000 m in Hubei and Sichuan). Our specimen resembles much with the *Mahonia hakeaeformis* hypotypes figured by BECKER (1969) in figure 4 of the plate XXVI. Anyway, this leaf is dissimilar to the other presented in the same group, by its small dimensions and marginal architecture. *B. coloradensis* is one of the most primitive forms (plesiomorphy) and makes problematic any assignment of this species to any extant section of the genus (SCHORN & WOLFE, 1989). HABLY et al. (2000) disagree with that and suggest that *B. coloradensis* and one of the four *Berberis* new species identified in Mexico (RAMÍREZ & CEVALLOS-FERRIZ, 2000), *B. lozanofolia*, represent foliage of barberries related to the extant South American section *Actinacanthae*. This very intriguing fossil record is for the first time described in the Tertiary of Europe and rise questionable problems about phytogeography of *Berberis*. When did it migrate from Eastern Asia (see GÜNER & DENK, 2012) and what kind of route did it follow? What is the relation between the occurrences of *B. coloradensis* from the Late Oligocene Creede flora (North America), *B. lozanofolia* RAMÍREZ & CEVALLOS-FERRIZ from the Oligocene of Puebla (Mexico), and our Middle Miocene specimen as closely related species? Moreover, it cannot be explained the morphologically conservative character of the taxa during the Oligocene epoch and up to the Middle Miocene. *B. coloradensis* in Ciocadia fossil assemblage may now be considered endemic until there are made further discoveries.

Order Urticales DUMORTIER 1829

Family Ulmaceae MIRBEL 1815 nom. cons.

Genus *Ulmus* LINNAEUS 1753

*Ulmus pyramidalis* GOEPPERT 1855

Pl. I, Figs. 4, 5, 6, 7, 8, 9, 10.

*Ulmus pyramidalis* GOEPPERT, GOEPPERT, 1855, p. 29, Pl. XIII, Figs. 10, 11, 12; Pl. XIV, Figs. 18, 19, 20.

*Ulmus Braunii* HEER, HEER, 1859, p. 181, Pl. CLI, Fig. 29, 31.

*Ulmus braunii* HEER, BARBU, 1936, p. 379, Text-Fig. 18.

*Ulmus pyramidalis* GOEPPERT, BŮŽEK, 1971, p. 56-57, Pl. XIX, Figs. 3-18.

Material: BCI.0184, BCI.0269, BCI.0302, BCI.0313, BCI.0328, BCI.0562c, BCI.0615.

Description. One-seeded (dry) fruits, with the pericarp modified into a narrow winglike structure (samara), membranous, adapted for airborne dispersal, ovate to elliptic and orbicular in shape, typically of 7.5-11.5 mm long and 4-5 mm wide. The base of the samara is concave, slightly asymmetric, stalked (1-4 mm long). The pedicels can often possess pieces of slightly thicker bracteoles (stipe) non-adherent to the base. Samaras conspicuously notched at the tip (beak-like tip or stigmatic cleft), and rarely with persistent lobed perianth, funnel-shaped pair of styles and bifurcate stigma. Seed (endocarp) compressed, distinct (of dark colour), elliptic to oval, 4 mm long and 1.5 mm wide, centrally or slightly toward base disposed on the samara; seed apex not reaching notch. Wings with strong axial vein laterally deflected from the stipe and connected to the seed, and radiant fine venation when visible.

Remarks. Leaves, fruits and pollen assignable to the Ulmaceae can be traced back to the Late Cretaceous, in North America, Europe, and Asia. Foliage and fruits of *Ulmus pyramidalis* are frequently cited in Europe from the Late Oligocene to the Pliocene time. Occurrence in the fossil floras of Romania: **Valea Jiului** (Zsil-Tales, Petroșani, Hunedoara County, Oligocene, leaves), **Valea Almașului** (Sălaj County, Oligocene, leaves), **Surdac** (Sălaj County, Late Oligocene, Chattian, leaves), **Muereasca de Sus** (Vâlcea County, Late Oligocene-Early Miocene, fruits), **Coruș** (Cluj County, Early Miocene=Late Egerian, Aquitanian, fruits and leaves), **Tihău** (Sălaj County, Early Miocene, Eggenburgian, Burdigalian, leaves), **Slătioara** (Vâlcea County, Miocene, fruits), **Valea Glâmboaca** (Vâlcea County, Badenian, fruits), **Căstău** (Orăștie, Hunedoara County, Early Badenian, fruits), **Pârlagele** (Mehedinți County, Late Badenian, leaves), **Porceni** (Gorj County, Sarmatian, leaves), **Râmestii** (Vâlcea County, Sarmatian, leaves), **Feleac** (Cluj County, Sarmatian, fruits), **Comănești** (Bașca County, Sarmatian, leaves), **Oaș** (Satu-Mare County, Early Sarmatian, leaves), **Daia** (Sibiu County, Early Sarmatian, Volhynian, leaves), **Săcădate** (Sibiu County, Early Sarmatian, Volhynian, leaves), **Morilor Valley** (Mehedinți County, Early Sarmatian, Volhynian-Early Basarabian, leaves and fruits), **Cavnic** (Maramureș County, Early Sarmatian, Volhynian-Early Basarabian, leaves), **Săcădate** (Sibiu County, Early Sarmatian, Volhynian-Early Basarabian, leaves), **Corni** (Neamț County, Middle Sarmatian, Basarabian, leaves), **Bunești** (Fălciu, Suceava County, Middle Sarmatian, leaves and woods, *Ulmoxylon kersonianum* STAROSTIN & TRELEA), **Tănăsești-Râmestii** (Vâlcea County, Late Sarmatian, Basarabian-Chersonian, leaves), **Hârșova** (Vaslui County, Late Sarmatian, leaves), **Păun** (Iași County, Late Sarmatian, Chersonian, leaves), **Valea Neagră de Criș** (Valea Crișului I & II, Bihor County, Pannonian B-C, leaves, GIVULESCU, 1997), **Beznea** (Delureni, Bihor County, Pannonian E, fruits), **Ibana Valley** (Mânzați, Vaslui County, Maeotian, leaves), **Turț** (Satu Mare County, Pontian, leaves), **Băița** (Maramureș County, Pontian, leaves), **Miclușoara** (Covasna County, Pontian, leaves), **Borsec** (Harghita County, Pontian, leaves), **Odești** (Maramureș County, Early Pontian, leaves), **Sărmășag** (Chieșd, Sălaj County, Early Pontian, leaves), **Crivina** (Mehedinți County, Early Pontian, leaves), **Batoți** (Mehedinți County, Early Pontian, leaves), **Cornițel** (Bihor County, Early Pontian, leaves, fruits), **Vișag** (Valea Nochi Mică, Timiș County, Middle

Pontian=Portaferrian, leaves), **Chiuzbaia** (Maramureş County, Late Pontian, leaves), **Cărbuneşti** (Prahova County, Late Pontian-Early Dacian, leaves), **Jilţ** (Jilţ Sud open-pit coal mine, Gorj County, Early Pliocene, Dacian, leaves), **Băceleş** (Dolj County, Late Pliocene, Middle Romanian, leaves), **Biborţeni** (Covasna County, Late Pliocene, leaves), and **Bodoş** (Covasna County, Late Pliocene-Pleistocene, leaves). There are signalled about 45 extant species of *Ulmus* (most speciose genus of the family) occurring widespread in the temperate and boreal regions of the Northern Hemisphere and extending into tropical America and Malesia (MANCHESTER, 1989). The centre of distribution seems to be central and northern Asia (KUBITZKI, 1993). According to CIOCĂRLAN (2009), in Romania there grow naturally only four *Ulmus* species: *U. laevis* PALLAS (syn. *U. effusa* WILLDENOW), *U. glabra* HUDSON (syn. *U. montana* STOKES; *U. scabra* MILLER), *U. procera* SALISBURY and *U. minor* MILLER (syn. *U. foliacea* GILIBERT ex C. K. SCHNEIDER; *U. campestris* auct non LINNAEUS; *U. carpinifolia* GLEDITSCH). Their ecology demonstrate high adaptation ability (BORLEA, 2004) concerning the soil humidity, preferring drained soils of typical forest steppe sites, but also can grow disseminated in easily flooded areas along river banks or in some meadows. *U. pyramidalis* GOEPPERT is commonly compared (BŮŽEK, 1971) with the modern *U. americana* LINNAEUS and *U. alata* MICHAUX, which are constituents of the riparian forests in the eastern and south-eastern USA. Apparently, the winged fruits from Ciocadia site match better with *U. americana* (section *Blepharocarpus* DUMORTIER) and probable not with *U. alata* of the section *Chaetoptelea* (LIEBMANN) C. SCHNEIDER. Currently, *U. pyramidalis* is considered a meso-hygrophyte member of intrazonal communities outside zonal communities, inhabiting as extrazonal element in gallery forests along high flood area of rivers, or sometimes in swampy conditions.

*Ulmus parschlugiana* KOVAR-EDER & KVAČEK 2004

Pl. I, Fig. 11.

*Ulmus braunii* HEER, BARBU, 1954, p. 41-42, Pl. VIII, Fig. 5.

*Ulmus parschlugiana* KOVAR-EDER & KVAČEK 2004; KOVAR-EDER, KVAČEK & STRÖBITZER-HERMANN, 2004, p. 67-68, Pl. VI, Figs. 23-26.

Material: BCI.0294.

Description. Larger single-seeded samaras, broadly oval, long stalked (stalk of 6 mm long). Endocarp distinct (of dark colour), with wing 10.5-15 mm long and 6.5-12 mm wide, elliptic to obovate in outline (4.5 mm long and 2.5 mm wide), styles persistent, 4 mm long. The seed body is typically oriented toward the base of samara and is surrounded by a broad membranous wing. Wings entire, very slightly undulate, usually do not have any distal stigmatic notch, or sometimes have a hint of a notch. The samara is flattened, where the seed is located, but not on the samara edges. Wing venation is clearly seen, composed of an axial vein laterally deflected from the stipe and connected to the seed, a surrounding strong marginal vein, and radiant reticulate fine venation from the seed, reticulate. Fruits are shortly stipitate, with stipe around 2 mm long.

Remarks. These kinds of endocarps with large wings are considered more evaluate because they may have improved the efficiency of fruit dispersal (DENK & DILLHOFF, 2005). The diversity of forms in the *Ulmus* fossils (leaves or samaras) is high in the Tertiary deposits of Europe, and is generally accepted the opinion of a greater specific differentiation in this genus. Large, broadly oval similar samaras are described in Romania from the **Morilor Valley** (Mehedinţi County, Early Sarmatian, PARASCHIV, 2004), **Porceni** (Gorj County, Sarmatian, BARBU, 1954), and **Valea Neagră de Criş** (Valea Crişului I & II, Bihor County, Pannonian B-C, GIVULESCU, 1997). Because in Ciocadia palaeoflora *Ulmus* leaves have not been yet described we cannot assign the fruits to the correspondent foliage. KOVAR-EDER et al. (2004) merged similar fruits of *U. parschlugiana* with the only one elm species which is foliage producer at Parschlug, *U. plurinervia* UNGER. *U. parschlugiana* is described from Parschlug (Styria, Austria) in deposits of Karpatian/Early Badenian age, based on the palaeofloristic composition. Fruits with broader wings and close morphology are dispersed by extant elm species like *U. americana* LINNAEUS, *U. minor* MILLER (an European polymorphic species), or the European elm with the widest natural range *U. glabra* HUDSON.

Genus *Zelkova* SPACH 1841

*Zelkova zelkovifolia* (UNGER 1843) BŮŽEK & KOTLABA in KOTLABA, 1963

Pl. I, Fig. 12.

*Zelkova (Planera) ungeri* KOVATS, BARBU, 1932, p. 13-14, Text-Fig. 9.

*Planera ungeri* KOVATS, BARBU, 1934, p. 126-127, Text-Fig. 31.

*Planera protokeaki* SAPORTA, BARBU, 1934, p. 127, Text-Fig. 32.

*Planera (Zelkova) Ungerii* KOVATS, BARBU, 1942, p. 130, Fig. 15.

Material: BCI.0129.

Description. Young branches with leaves having inconspicuously pedicelate axillary fruits in anatomical connection. Leaves simple, slightly asymmetric, oblong ovate to elliptic, 25-31 mm long and 10-13 mm wide, alternate on the shoot, with symmetric or slightly asymmetrical base, rounded, apex shortly acute, slightly bent sideways,

gradually narrowed, and coarse simple-serrated to crenulated margins (with “*Zelkova*-like teeth” or “wavy-toothed”), regularly spaced. Apical side of teeth short, acuminate, basal side convex, angular shape of the sinus, and the tooth apex is simple. Petioles are strong, 1.5-3 mm long. Venation simple craspedodromous, with midrib strong, straight, distally bent. Secondary veins conspicuous, parallel, alternate, originating at 45-70°, seven to nine pairs of veins, distally curved upward, and innervating characteristically the apices of teeth (their apical branches terminate in tooth apex forming an angle of about 40° with primary vein). Tertiary veins and venation of higher order are not obvious. The fruits are dry, nut-like drupe, not winged, subglobose to carinate (after compression are obvious bilobed), with stigmas beak-shaped, oblique attached to the leaf insertion on the shoot, approximately 3 mm in diameter.

Remarks. The specimen, which is a young (current-season) fertile shoot with conspicuously zig-zag architecture, preserves three fruits, solitary, in the axils of upper leaves. This is a rare case with leaves that occur together with fruits in natural state, because *Zelkova* usually grew at some distance from the sedimentary basins (DENK & DILLHOFF 2005). Leaf fossils attributed to *Zelkova* are particularly common in Eurasia from the Eocene-Oligocene to the Plio-Pleistocene. This taxon is not very uniform as regards the general morphology of leaves (continuous transition *Ulmus-Zelkova*, comprising also common-size leaves, see BŮŽEK, 1971) but, in our case, the co-occurrence with characteristic fruits excludes any confusion. *Z. zelkovifolia* is widespread in the Tertiary floras of Romania: **Valea Almașului** (Sălaj County, Oligocene, leaves and fossil woods, *Zelkovoxylon dacicum* PETRESCU), **Slătioara** (Vâlcea County, Miocene), **Coruș** (Cluj County, Early Miocene=Late Egerian, Aquitanian), **Tihău** (Sălaj County, Early Miocene), **Pătărlagele** (Buzău County, Late Badenian), **Baia Sprie** (Maramureș County, Sarmatian), **Racșa** (Satu-Mare County, Early Sarmatian), **Deva** (Hunedoara County, Early Sarmatian, Early Volhynian), **Morilor Valley** (Mehedinți County, Early Sarmatian), **Tâmpa** (Hunedoara County, Early Sarmatian, as a fertile branch with three leaves and two fruits attached axillary), **Oaș** (Satu-Mare County, Early Sarmatian), **Pârteștii de Sus** (Suceava County, Early Sarmatian, Volhynian), **Corni** (Neamț County, Middle Sarmatian), **Bunești** (Fălciu, Suceava County, Middle Sarmatian), **Tănăsești-Râmești** (Vâlcea County, Late Sarmatian), **Păun** (Iași County, Late Sarmatian), **Valea Neagră de Criș** (Valea Crișului I & II, Bihor County, Pannonian B-C, leaves with fruits), **Vadu Crișului** (Bihor County, Early Pannonian), **Beznea** (Delureni, Bihor County, Pannonian E), **Tirol** (Caraș-Severin County, Pannonian G), **Ibana Valley** (Mânzați, Vaslui County, Maeotian), **Sinersig** (Timiș County, Pontian), **Miclușoara** (Covasna County, Pontian), **Herculian** (Covasna County, Pontian), **Hărman** (Brașov County, Pontian), **Borsec** (Harghita County, Pontian), **Batoți** (Mehedinți County, Early Pontian), **Cornițel** (Bihor County, Early Pontian), **Vișag** (Valea Nochi Mică, Timiș County, Middle Pontian=Portaferrian), **Chiuzbaia** (Maramureș County, Late Pontian), **Cerna** (open-pit coal mine, Vâlcea County, Early Pliocene, Dacian), **Baraolt** (“Grădina cu mei”, Covasna County, Late Pliocene, Early Villafranchian), **Biborțeni** (Bibartfalva, Covasna County, Late Pliocene), **Doboșeni** (Covasna County, Late Pliocene-Pleistocene), and **Bodoș** (Covasna County, Late Pliocene-Pleistocene). There are six extant species of *Zelkova*, two Mediterranean: *Z. abelicea*, originating from Crete and *Z. sicula* from south-eastern Sicily (NAKAGAWA et al., 1998), one Caucasian (Turkey, Georgia, Armenia, Azerbaijan, Iran) *Z. carpinifolia* (KVAVADZE & CONNOR, 2005), and three east Asian: the Chinese species *Z. schneideriana* and *Z. sinica* and the Japanese *Z. serrata* (DENK & GRIMM, 2005). *Z. zelkovifolia* can be compared with the recent species *Z. carpinifolia* (PALLAS) K. KOCH (= *Z. crenata* SPACH) from relic deciduous and mesophytic forests formations of Colchis (Caucasus, Kaçkar, and Alborz mountains). This mesophilous tree flourishes on moist well-drained soils and warm conditions, and do not show resilience to frost or tolerance to water logging or swampy conditions. *Z. zelkovifolia* is included in the (Warm-) temperate Broad leaved Deciduous Forest as a zonal vegetation formation (KVAČEK et al., 2006). It represents the so called “edaphic vegetation” because it probably lived on more humid soils along shallow streams that entered the sea, but also it was an important component of the mesophytic open woodlands.

## RESULTS AND DISCUSSIONS

Taxa like *Berberis coloradensis*, *Ulmus pyramidalis* and *U. parschlugiana* show a combination of long stratigraphical ranges and wide geographic range, and presumably can demonstrate a considerable tolerance to changing environmental factors (TANAI & WOLFE, 1977). This study demonstrates that the occurrence of *Ulmus* fruits in the fossil assemblages is scarce and that its leaves are best preserved in sediments. Only 9 fossil floras from the total of 44 with *U. pyramidalis* remains are documented by fruits (and leaves). There is a large amount of *Ulmus* samaras in the Ciocadia palaeoflora and this can be explained by a characteristic dominance of this taxon. Because *Zelkova* twigs are poorly dispersed over long distances, the presence in the Ciocadia assemblage likely indicates its local origin and a significant participation together with *Ulmus* in palaeoforests.

## CONCLUSIONS

This is the second part of a series of contributions, which will clarify the floristic and vegetation aspects of the Middle Miocene flora of Ciocadia (Oltenia Province). In this paper we describe well-preserved fossils of leaves attached to fruits belonging to *Zelkova*, leaves assignable to two different species of *Berberis*, and isolated fruits of *Ulmus*. We also provide the necessary data about the general occurrence in Romania of mentioned taxa and thus contribute to a better understanding of the evolution in Berberidaceae and Ulmaceae.

## ACKNOWLEDGEMENTS

I am grateful to Prof. Zlatko Kvaček from Charles University in Prague, for helpful discussions and useful suggestions regarding the fossil materials from Ciocadia site. This work was supported by the grants ID\_442/2007 and ID\_584/2008, financed by CNCSIS (the National University Research Council) and also by the grant POSDRU/107/1.5/S/80765 of the European Social Fund.

## REFERENCES

- AXELROD D. I. 1987. *The Late Oligocene Creede flora, Colorado*. University of California Publications in Geological Sciences. University of California Press. Berkeley. **130**: 1-236.
- BARBU I. Z. 1954. *Flora fosilă din Terțiarul Olteniei*. Anuarul Comitetului Geologic. Institutul Geologic. București. **27**: 5-76.
- BECKER H. F. 1969. *Fossil plants of the Tertiary Beaverhead Basins in Southwestern Montana*. Palaeontographica. Schweizerbart'sche Verlagsbuchhandlung (Nägele u. Obermiller). Stuttgart. **127**(1-6): 1-142.
- BOHN-HAVAS M., LANTOS M., SELMECZI I. 2004. *Biostratigraphic studies and correlation of Tertiary planktonic gastropods (Pteropods) from Hungary*. Acta Palaeontologica Romaniaiae. Edit. Supergraph. Cluj-Napoca. **4**: 37-43.
- BORLEA G. F. 2004. *Ecology of elms in Romania*. Investigación Agraria: Sistemas y recursos forestales. Ed. by Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria. Madrid. **13**(1): 29-35.
- BŮŽEK Č. 1971. *Tertiary flora from the Northern part of the Pětipsy area (North-Bohemian Basin)*. Rozpravy Ústředního Ústavu Geologického. Vydal Ústřední Ústav Geologický. Praha. **36**: 1-120.
- CIOCĂRLAN V. 2009. *Flora ilustrată a României. Pteridophyta et Spermatophyta*. Edit. Ceres. București. 1141 pp.
- DENK T. & DILLHOFF R. M. 2005. *Ulmus leaves and fruits from the Early-Middle Eocene of northwestern North America: systematics and implications for character evolution within Ulmaceae*. Canadian Journal of Botany. National Research Council Press. Ottawa. **83**: 1663-1681.
- DENK T. & GRIMM G. W. 2005. *Phylogeny and biogeography of Zelkova (Ulmaceae sensu stricto) as inferred from leaf morphology, ITS sequence data and the fossil record*. Botanical Journal of the Linnean Society. Edit. John Wiley & Sons Inc. London. **147**: 129-157.
- GIVULESCU R. 1990. *Flora fosilă a Miocenului superior de la Chiuzbaia (Judetul Maramureș)*. Edit. Academiei Române. București. 238 pp.
- GIVULESCU R. 1997. *Istoria pădurilor fosile din Terțiarul Transilvaniei, Banatului, Crișanei și Maramureșului*. Edit. Carpatica. Cluj-Napoca. 172 pp.
- GIVULESCU R. 2002. *About the possible presence of Berberis kymeana (Unger) Kvaček et Erdei in the Early Pannonian flora from Valea Crișului (Bihor County - Romania)*. Contribuții Botanice. Grădina Botanică „Alexandru Borza”. Cluj-Napoca. **37**: 289-291.
- GOEPPERT H. R. 1855. *Die Tertiäre flora von Schosnitz in Schlesien*. Heyn'sche Buchhandlung (E. Remer.). Görlitz. 52 pp.
- GÜNER T. H. & DENK T. 2012. *The genus Mahonia in the Miocene of Turkey: Taxonomy and biogeographic implications*. Review of Palaeobotany and Palynology. Edit. Elsevier. Amsterdam. **175**: 32-46.
- HABLY L., KVAČEK Z., MANCHESTER S. R. 2000. *Shared taxa of land plants in the Oligocene of Europe and North America in context of Holarctic phytogeography*. Acta Universitatis Carolinae. Geologica. Univerzita Karlova. Praha. **44**: 59-74.
- HEER O. 1859. *Flora Tertiaria Helvetiae*. Dritter Band, Die Gamopetalen und Polypetalen Dicotyledonen. Verlag der lithographischen Anstalt von Wurster & Comp. Winterthur. 378 pp.
- HICKEY L. J. 1979. *A revised classification of the architecture of Dicotyledonous leaves*. In: Metcalfe R. C. & Chalk L. (Eds.): Anatomy of the Dicotyledons, Systematic anatomy of leaf and stem, with a brief history of the subject. Clarendon Press. Oxford. **1**: 25-39.
- HUICĂ I. V. 1994. *Recherches géologiques sur le Badénien et la Sarmatien de la région de Ciocadia-Săcel (l'anticlinal Săcel-Tg. Jiu, Dépression Gétique)*. The Miocene from the Transylvanian Basin, Romania. Edit. Carpatica. Cluj-Napoca: 163-170.
- IORGULESCU T. 1952. *Elemente de micropaleontologie aplicată*. Edit. Tehnică. București. 245 pp.
- KOVAR-EDER J., KVAČEK Z., STRÖBITZER-HERMANN M. 2004. *The Miocene flora of Parschlug (Styria, Austria) – revision and synthesis*. Annalen des Naturhistorischen Museums in Wien. Selbstverlag Naturhistorisches Museum. Wien. **105 A**: 45-160.
- KUBITZKI K. 1993. *Flowering plants. Dicotyledons. Magnoliid, Hamamelid and Caryophyllid Families*. In: Kubitzki K., Rohwer J. G., Bittrich V. (Eds.). The families and genera of vascular plants. Springer Verlag. Berlin. 653 pp.
- KVAČEK Z., DVOŘÁK Z., MACH K., SAKALA J. 2004. *Třetihorní rostliny severočeské hnědouhelné pánve*. Severočeské doly a. s. Chomutov Granit, s. r. o. Praha. 160 pp.
- KVAČEK Z., KOVÁČ M., KOVAR-EDER J., DOLÁKOVÁ N., JECHOREK H., PARASCHIV V., KOVÁČOVÁ M., SLIVA L. 2006. *Miocene evolution of landscape and vegetation in the Central Paratethys*. Geologica Carpathica. Publishing House of the Slovak Academy of Sciences. Bratislava. **57**(4): 295-310.
- KVAVADZE E. V. & CONNOR S. E. 2005. *Zelkova carpinifolia (PALLAS) K. KOCH in Holocene sediments of Georgia – an indicator of climatic optima*. Review of Palaeobotany and Palynology. Elsevier. Amsterdam. **133**(1-2): 69-89.


- LI Y.-L., KVAČEK Z., FERGUSON D.K., WANG Y.-F., LI C.-S., YANG J., YING T.-S., ABLAEV A.G., LIU H.-M. 2010. *The fossil record of Berberis (Berberidaceae) from the Palaeocene of NE China and interpretations of the evolution and phytogeography of the genus*. Review of Palaeobotany and Palynology. Elsevier. Amsterdam. **160**(1-2): 10-31.
- MANCHESTER S. R. 1989. *Systematics and fossil history of the Ulmaceae*. Evolution, systematics, and fossil history of the Hamamelidae. Higher Hamamelidae. The Systematics Association Special Volume. Clarendon Press. Oxford. **40B**: 221-251.
- MARINESCU F. 1969. *Precizări stratigrafice privind Sarmatianul și Meoțianul din Nordul Olteniei*. Dări de Seamă ale Ședințelor. Institutul Geologic al României. București. **54**(3): 153-162.
- NAKAGAWA T., GARFI G., REILLE M., VERLAQUE R. 1998. *Pollen morphology of Zelkova sicula (Ulmaceae), a recently discovered relic species of the European Tertiary flora: description, chromosomal relevance, and palaeobotanical significance*. Review of Palaeobotany and Palynology. Elsevier. Amsterdam. **100**(1-2): 27-37.
- PALAMAREV E. H. & PETKOVA A. S. 1987. *La Macroflore du Sarmatien*. In: Palamarev E. H. & Petkova A. S. Les fossiles de Bulgarie. Academie Bulgare des Sciences. Sofia. 276 pp.
- PARASCHIV V. 2004. *The fossil flora of the Morilor Valley, South-Western Dacian Basin, Romania*. Acta Palaeontologica Romaniae. Edit. Supergraph. Cluj-Napoca. **4**: 315-330.
- PARASCHIV V. 2006. *Flora Sarmatiană din Oltenia*. Unpublished PhD Thesis. Universitatea din București. 263 pp.
- PARASCHIV V. 2008a. *New data concerning the fossil flora of the Ciocadia Valley, Western Dacian Basin, Romania*. Acta Horti Botanici Bucurestiensis. Universitatea din București. Grădina Botanică „Dimitrie Brândza“. București. **35**: 36-42.
- PARASCHIV V. 2008b. *New Sarmatian plant macroremains from Oltenia region (Romania), Part I*. Acta Palaeontologica Romaniae. Universitatea „Alexandru Ioan Cuza“ Iași. **6**: 279-286.
- PARASCHIV V. & SEBE O.-G. 2011. *New contributions to the Ciocadia Middle Miocene flora (part one)*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **27**: 183-190.
- PETRESCU I. & DRAGASTAN O. 1981. *Plante fosile, introduce în paleobotanică*. Edit. Dacia. Cluj-Napoca. 471 pp.
- PETRESCU I., GIVULESCU R., BARBU O. 1997. *Macro- și microflora oligocenă de la Cornești-Aghireș, România*. Edit. Carpatica. Cluj-Napoca. 218 pp.
- POPESCU G. 1953. *Cercetări geologice în regiunea Ciocadia-Pițicu-Baia de Fier (Depresiunea Getică)*. Dări de Seamă ale Ședințelor. Institutul Geologic al României. București. **39**: 15-25.
- RAMÍREZ J. L. & CEVALLOS-FERRIZ S. R. S. 2000. *Leaves of Berberidaceae (Berberis and Mahonia) from Oligocene sediments, near Tepexi de Rodríguez, Puebla*. Review of Palaeobotany and Palynology. Elsevier. Amsterdam. **110**(3-4): 247-257.
- SĂVULESCU T., BELDIE AL., BUIA AL., GRINȚESCU G., GRINȚESCU I., GUȘULEAC M., NYÁRÁDY A., NYÁRÁDY E. I., RĂVĂRUȚ M., ȘERBĂNESCU I., ȚOPA E., ZAHARIADI C. 1955. *Flora R. P. Române*. Edit. Academiei R. P. R. București. 662 pp.
- SCHOPF T. J. M. 1980. *Paleoceanography*. Harvard University Press. Cambridge. 341 pp.
- SCHORN H. E. & WOLFE J. A. 1989. *Taxonomic revision of the Spermatopsida of the Oligocene Creede flora, Southern Colorado*. U. S. Department of the Interior. Geological Survey. Open-File Report. Washington. **89-433**: 1-95.
- TAKHTAJAN A. 2009. *Flowering plants*. Second edition. Springer Science. Amsterdam. 871 pp.
- TANAI T. & WOLFE J. A. 1977. *Revisions of Ulmus and Zelkova in the middle and late Tertiary of western North America*. Geological Survey Professional Paper. U. S. Government Printing Office. Washington. **1026**: 1-23.
- TUDOR M. 1955. *Stratigrafia și fauna depozitelor Tortoniene și Sarmatiene dintre Jiu și Olteț*. Edit. Academiei Române. București. 118 pp.
- ȚICLEANU N. 1984. *Contributions to the study of the Upper Badenian flora from Romania*. Dări de Seamă ale Ședințelor. Institutul Geologic al României. București. **68**(3): 135-150.
- ȚICLEANU N. & MICU M. 1979. *Flore Sarmatiene de Corni (District de Neamt)*. Dări de Seamă ale Ședințelor. Institutul de Geologie și Geofizică. București. **64**(3): 399-414.
- ȚICLEANU N. & GIVULESCU R. 1978. *Contributions to the knowledge of the Upper Egerian palaeoflora - fossil flora of Coruș II, Cluj District*. Courier Forschungsinstitut Senckenberg. E. Schweizerbart'sche Verlagsbuchhandlung (Nägele u. Obermiller). Frankfurt am Main. **30**: 133-150.
- VELCESCU M. 1997. *Pteropod assemblage in the Miocene sediments of the Baia Mare region*. Acta Palaeontologica Romaniae. Edit. Eficient. București. **1**: 161-164.
- WING S., ASH A., ELLIS B., HICKEY L. J., JOHNSON K., WILF P. 1999. *Manual of leaf architecture. Morphological description and categorization of Dicotyledonous and net-veined Monocotyledonous Angiosperms*. Smithsonian Institution. Washington, DC: 1-67.

Paraschiv Valentin

Geological Institute of Romania, National Museum of Geology,  
Kiseleff Ave, No. 2, Sect. 1, 011345, Bucharest, Romania.  
E-mail: paleoval@yahoo.com

Received: March 05, 2013

Accepted: May 30, 2013



Scale bar in millimeters

**PLATE I**

Fig. 1, 2 - *Berberis* sp.

Fig. 3 - *Berberis coloradensis* AXELROD 1987

Fig. 4-10 - *Ulmus pyramidalis* GOEPPERT 1855

Fig. 11 - *Ulmus parschlugiana* KOVAR-EDER & Z. KVAČEK 2004

Fig. 12 - *Zelkova zelkovifolia* (UNGER 1843) BŮŽEK & KOTLABA in KOTLABA 1963