

IN VITRO MULTIPLICATION OF *Acer platanoides* L.**Vladimir TOMOV, Ivan ILIEV**

Abstract. Mature individuals of “Faassens Black” and “Drummondii” were used as stock plants. Hardwood nodal and apical segments, (harvested in September), softwood nodal and apical segments (harvested in May), and dormant buds (harvested in the end of April and November) were used as initial explants. Dormant buds are the most suitable explants for disinfection and culture establishment on PGRs-free WPM. The concentration of kinetin does not affect the length of the shoots within the cultivar, but in comparison with “Drummondii” they were statistically longer in “Faassens Black” on WPM with 0.5 mg l⁻¹ kinetin (14.5 ± 3.5 mm). However, axillary shoots formation was not induced in both treatments and cultivars. These shoots in “Faassens Black i” reached maximal length (39.5 ± 11.0 mm) after the 8th subculture on WPM supplemented with 1.0 mg l⁻¹ BAP and 0.1 mg l⁻¹ IBA. In “Drummondii” only single shoots elongated but all of them gradually died and no viable shoots were found after the 8th subculture. However, the frequency of the induced axillary shoots after cultivation for 180 days was low and was noticed on single explants in “Faassens Black”, only (0.65 ± 0.17). In juvenile explants, the multiplication rate and the length of the axillary shoots were low and were not affected by the genotype and subcultures. Maximal number (1.9 ± 0.4) and length (8.4 ± 0.6 mm) of axillary shoots was induced on the medium with 1.0 mg l⁻¹ BAP and 1.0 mg l⁻¹ GA₃. The combination of these PGRs provoked statistically significant additional formation of axillary shoots but their length remained less than 5 mm in the period of cultivation. The type of cytokinin does not affect the elongation of the initial explant within the genotype. The quantity of induced axillary shoots was small and their elongation was very slow. A higher number of axillary shoots was received on WPM with 2 mg l⁻¹ Kinetin (from 0.25 ± 0.16 to 2.13 ± 0.58). Probably, due to the apical dominance of the initial explant, the formation of axillary shoots, and their elongation is limited. Also, different responses to cytokinins could be attributed to the reaction of different genotypes.

Keywords: “Drummondii”, “Faassens Black”, Norway maple, propagation, subculture, tissue cultures.

Rezumat. Înmulțirea *in vitro* la *Acer platanoides* L. Exemplare mature de „Faassens Black” și „Drummondii” au fost folosite ca plante stoc. Ca explante inițiale au fost folosite segmente nodale și apicale de lemn tare (recoltate în luna septembrie), segmente nodale și apicale de lemn moale (recoltate în mai) și muguri latenți (recoltați la sfârșitul lunii aprilie și noiembrie). Mugurii latenți sunt cei mai potriviti ca explante pentru dezinfecție și stabilirea culturii pe WPM fără PRG. Concentrația de kinetină nu a afectat lungimea lăstarilor în cazul aceluiași soi, dar în comparație cu „Drummondii”, aceștia au fost statistic mai lungi în cazul „Faassens Black” pe WPM cu 0,5 mg l⁻¹ kinetină (14.5 ± 3.5 mm). Cu toate acestea, formarea de muguri axilari nu a fost indușă în cazul ambelor tratamente și soiuri. Acești lăstari de „Faassens Black” au atins lungimea maximă (39.5 ± 11.0 mm) după subcultura a opta pe WPM suplimentat cu 1,0 mg l⁻¹ BAP și 0,1 mg l⁻¹ IBA. La „Drummondii”, numai lăstari unici s-au alungit, dar aceștia au murit treptat și nu au fost depistați muguri viabili după o opta subcultura. Cu toate acestea, frecvența lăstarilor induși axial după cultivare timp de 180 de zile a fost scăzută și a fost observată doar la explantele unice de „Faassens Black” (0.65 ± 0.17). La explantele juvine, rata de multiplicare și lungimea lăstarilor axilari au fost reduse și nu au fost afectate de genotip și subcultură. Numărul maxim (1,9 ± 0,4) și lungimea maximă (8,4 ± 0,6 mm) a mugurilor axilari a fost indușă pe un mediu cu 1,0 mg l⁻¹ BAP și 1,0 mg l⁻¹ GA₃. Combinarea acestor PGR a determinat formarea adițională de lăstari axiali, semnificativă din punct de vedere statistic, dar lungimea acestora nu a depășit 5 mm în perioada de cultivare. Tipul de citokinină nu afectează alungirea explantului inițial în cadrul aceluiași genotip. Cantitatea generată de lăstari axiali a fost mică și alungirea lor a fost foarte lentă. Un număr mai mare de lăstari axilari a fost determinat în cazul WPM cu 2 mg l⁻¹ kinetină (de la 0.25 ± 0.16 - 2.13 ± 0.58). Probabil, datorită dominației apicale a explantului inițial, formarea de lăstari axilari și alungirea lor este limitată. De asemenea, răspunsurile diferite la citokinine ar putea fi atribuite reacțiilor diferitelor genotipuri.

Cuvinte cheie: „Drummondii”, „Faassens Black”, arțar norvegian, propagare, subcultură, culturi de țesuturi.

INTRODUCTION

Acer platanoides L. is a valuable ornamental tree with a lot of ornamental cultivars (ARTJUSHENKO et al., 1958; KOLESNIKOV, 1958; SANTAMOUR et al., 1982; KRÜSSMAN, 1984; DIRR, 1998). It is readily propagated from seeds, but as seedlings show a wide variability in phenotypic characteristics, vegetatively propagated cultivars are preferable in urban areas. Norway maple cuttings are considered difficult to root (CHONG & DAIGNEAULT, 1986; MAYNARD & BASSUK, 1987) and depend on the age of the stock plant and etiolation of the shoots (MAYNARD & BASSUK, 1987). For this reason most of the cultivars are grafted onto seedlings origin rootstock. Furthermore, the production of large quantities of grafts is limited by the season, the period for rootstock production and the success depends on the method of grafting and climatic conditions in the field (HOWARD et al., 1974; KRÜSSMAN, 1984; DIRR & HEUSER, 1987; HOWARD, 1993; HOWARD & OAKLEY, 1997).

In vitro propagation can provide alternative of the conventional methods for cloning of valuable ornamental genotypes (BONGA, 1982; HARTMANN et al., 2002; ILIEV & BORISOWA, 2002). Studies on the *in vitro* propagation of the Norway maple have focused on wild types (ĐURKOVIĆ, 1996; LINDEM & RIIKONEN, 2006) or ornamental “Crimson Sentry” (CHENG, 1978; LATTIER et al., 2012, 2013), “Crimson King” (MARKS & SIMSON, 1990, 1994), and “Jirka” (ŠEDIVÁ, 2009). However, some of the publications did not report the age of the stock plant and multiplication did not accomplish (CONCIOU et al., 2010; ROVINĀ et al., 2010)

This research was aimed at determining a suitable disinfection, factors affecting culture establishment, and multiplication of mature "Faassens Black" and "Drummondii" and to compare their reaction *in vitro* with juvenile *Acer platanoides* L. seedlings.

MATERIAL AND METHOD

Factors affecting the cultivation of mature genotypes

Plant material and disinfection of the initial explants

Individuals of "Faassens Black" and "Drummondii" were used as stock plants 8 years after their grafting. Hardwood nodal and apical segments (≈ 3 cm) were harvested in September and were disinfected consecutively by 72% ethanol for 2 min and then by 0.2% $HgCl_2$ for 15 min (variant 3). Softwood nodal and apical segments (≈ 3 cm) were harvested in May and were disinfected by 0.1% $HgCl_2$ for 3 min (variant 2). Dormant buds were harvested in the end of April and November and were disinfected consecutively by 72% ethanol for 2 min and then by 0.2% $HgCl_2$ for 8 min (variant 1). After being rinsed 3 times in sterile distilled water, the ends (≈ 2 mm) of apical and nodal segments were aseptically removed.

Culture establishment

Hardwood nodal and apical segments were cultured on plant growth regulators (PGRs)-free WPM (LLOYD & McCOWN 1980) or WPM supplemented with 1.0 mg l⁻¹ 6-Furfurilaminopurine (Kinetin), and the buds were cultured on PGRs-free WPM. Softwood nodal and apical segments were cultured on PGRs-free WPM or WPM supplemented with 6-benzylaminopurine (BAP), (6-[4-Hydroxy-3-methyl-but-2-enylamino]purine (Zeatin) or Kinetin in concentration 1.0 mg l⁻¹. After two days of cultivation, the explants were subcultured on the same medium for the elimination of the separated phenols (Fig. 1).




Figure 1. Released phenols on the second day after the cultivation of softwood explants (original).

Each treatment contained three repetitions, and 15 explants were used per replication. The explants with developed buds and elongated shoots as well as dead explants were evaluated after 2 weeks.

Factors affecting the elongation and multiplication of the shoots

After 2 weeks on PGRs-free WPM, the buds with removed scales harvest in April from "Faassens Black" and "Drummondii" were cultured on WPM supplemented with 0.5 and 1.0 mg l⁻¹ kinetin for elongation and multiplication. Three replications, each containing 6-7 were used for each variant of the medium. After 14 days, the explants were subcultured on the same variant of the medium and the results were evaluated after 28 days from the beginning of the experiment.

The elongated shoot tips (~ 2 cm) were cultured on WPM supplemented with 1.0 mg l⁻¹ BAP and 0.1 mg l⁻¹ indole-3-butryic acid (IBA). The effect of 10 subcultures (each with a duration of 14 days) was studied on the reaction of initial explants and axillary shoots formation. The results were evaluated after each 2nd subculture and the formed axillary shoots were isolated and cultured on the same medium. Finally, the isolated shoots were subcultured after 20 days, and the results were evaluated 20 days later. For each cultivar at least 14 initial explants were used at the beginning of the experiment.

Factors affecting the cultivation of juvenile genotypes

Effect of the genotype on the multiplication rate

To study the effect of genotype, the formed axillary shoots from 8 seedlings were cultured on WPM supplemented with 1.0 mg l⁻¹ kinetin and 1.0 mg l⁻¹ Gibberellic acid (GA₃). Due to the limiting initial material, 3 to 11 axillary shoots were used per genotype. The explants were subcultured after 14 days and the results were evaluated 14 days later.

Effect of subculture and genotype on the axillary shoots formation

To study the effect of genotype and subculture axillary shoots of three clones (2, 5, and 8) were cultured on WPM enriched with 1.0 mg l⁻¹ BAP and 0.1 mg l⁻¹ indole-3-butryic acid (IBA). The formed axillary shoots were subcultured 5 times and each subculture was with duration of 14 days. The number of formed axillary shoots was evaluated after each subculture.

Effect of PGRs on the axillary shoots formation

Axillary shoots (5-8 mm) from genotype 2 with small part of the callus (1-2 mm) on the base (Fig. 2) were cultured on WPM supplemented with 3.0 mg l⁻¹ BAP and 0.1 mg l⁻¹ IBA, 1.0 mg l⁻¹ BAP and 0.1 mg l⁻¹ IBA, 1.0 mg l⁻¹ BAP and 1.0 mg l⁻¹ GA₃, and 1.0 mg l⁻¹ N⁶-(2-Isopentenyl)adenine (2iP). Each treatment contained 3 repetitions and each of them - 6 or 7 explants. The explants were subcultured after 20 days on the same variants of the medium and the results were evaluated on the 40th day from the beginning of the experiment.



Figure 2. Axillary shoots with little callus on the base, used as initial explants (original).

Axillary shoots (10-15 mm), were used as initial explants (Fig. 3). They were cultivated on WPM medium supplemented with 1 mg l⁻¹ BAP, mg l⁻¹ 2iP, and 2.0 mg l⁻¹ Kinetin. Each treatment contained 8 explants. The shoots were transferred on the 20th day on the same variants of medium, after removing only the necrotic callus or leaves. Number and length of initial elongated and new formed axillary shoots were evaluated after 40 days. No mean length was calculated when the new formed shoots were less than 5 mm.



Figure 3. Axillary shoots, used as initial explants (original).

Conditions of the cultivation

Each variant of the media contained 7 g l⁻¹ agar (Sigma) and pH was adjusted to 5.6 before autoclaving (under pressure of 118 kPa and 120°C for 20 min). The cultures were grown in a cultivation chamber at 23 ± 1°C with 16 hrs of cool white fluorescent light at a photosynthetic photon flux density of 40 µmol m⁻² s⁻¹, daily.

Statistical analysis

The results were analysed by One-Way ANOVA followed by a post hoc LSD test at $p < 0.05$, using SPSS 20.0 for Windows. Percentage values were transformed using arcsine square root (\sqrt{P}) (COMPTON, 1994) to normalize error distribution prior variance analysis.

RESULTS AND DISCUSSION

Factors affecting the cultivation of mature genotypes

Disinfection of the explants and culture establishment

The results from disinfection experiments demonstrated that the highest statistically significant level of exogenous contamination was observed on softwood explants from both cultivars harvested in May. It could be due to the lower concentration of $HgCl_2$ and shorter duration of disinfection necessary for this type of explants. The level of exogenous contamination in other types of explants was low in the stage of their dormancy and significant differences between them and the time of their collection was not observed in both investigated cultivars. The highest statistically significant level of endogenous contamination in both cultivars was found in hardwood segments, collected in September. The level of endogenous contamination in other type of explants was low and significant differences between them and time of their collection was not observed in both investigated cultivars. The highest level of aseptic explants in both cultivars was obtained when buds were collected in April and November in the stage of their dormancy (Table 1). All our explants were resistant to the action of $HgCl_2$ i.e. they were not damaged from the treatment and were viable. Softwood explants and buds demonstrated development (Fig. 4). The elongation of the shoot was induced only from the dormant buds with removed scales, harvested in the beginning of the spring and cultivated on PGRs-free WPM. However, the percentage of the buds with elongated shoots was low in both cultivars (24.4 ± 8.0 and 11.1 ± 4.4 , resp.) and the length of the induced shoots reached not more than 5 mm in the 14 days of cultivation (Table 2, Fig. 5).



Figure 4. Softwood explants from *Acer platanoides* ‘Drummondii’ and ‘Faassens Black’, after 14 days of cultivation. No shoot elongation noticed (original).




Figure 5. Non elongation and elongation of buds from *Acer platanoides* ‘Faassens Black’.

The initial explants have been treated with 70% ethyl alcohol for 1 min before applying the disinfection agent (LINDEN & RIIKONEN, 2006). NaClO (5.0-6.15%) has been used for 15-20 min for disinfection of actively growing shoots (PREECE et al., 1991; LATTIER et al., 2013) or dormant segments (CONCIOU et al., 2010). Dormant apical and axillary buds were disinfected in 0.5% NaClO for 20 min (LINDEN & RIIKONEN, 2006). However, in some reports, the physiological condition of the initial explant is not indicated (ŠEDIVA, 2009; ROVINĂ et al., 2010). It is well known that HgCl₂ is the most effective disinfectant and for softwood explants it is used in 0.1% for 1-3 min and for hardwood explants in 0.2% for 8-10 min (ILIEV et al., 2010). Actively growing explants were disinfected by 0.5% HgCl₂ for 5 min followed by 5% NaClO for 15 min (MARKS & SIMPSON, 1990). However, in all these publications the percentage of aseptic explants is not reported. ROHR & HANUS (1987) reported that the receiving of aseptic softwood segments in *Acer pseudoplatanus* is very difficult after disinfection with 7.5% calcium hypochlorite for 15-20 min. ĐURKOVIĆ (2003) reported that dormant buds with removed scales have been successfully disinfected with 0.1% HgCl₂ for 2-5 min and the contamination rate of juvenile buds was 39.6% and of mature twice less (ĐURKOVIĆ 2003). Our results are in agreement with these findings and showed that dormant buds, isolated in April are the most suitable explants for disinfection (Table 1) and culture establishment (Table 2).

Table 1. Factors affecting the disinfection of the initial explants.

Period of harvesting	Variant of disinfection	‘Faassens Black’			‘Drummondii’		
		Exogenous contamination (%)	Endogenous contamination (%)	Aseptic explants (%)	Exogenous contamination (%)	Endogenous contamination (%)	Aseptic explants (%)
Hardwood segments							
September	3	13.3 ± 6.7 a	77.8 ± 8.0 c	8.9 ± 8.9 a	8.9 ± 2.2 a	73.3 ± 3.8 c	17.8 ± 4.5 ab
Softwood segments							
May	2	51.7 ± 5.8 b	3.9 ± 4.0 ab	34.4 ± 5.1 b	45.6 ± 6.8 b	23.9 ± 4.5 b	30.5 ± 3.7 b
Buds							
April	1	4.5 ± 2.2 a	3.3 ± 3.8 ab	82.2 ± 2.2 c	6.7 ± 6.7 a	8.9 ± 4.4 ab	84.5 ± 9.7 c
November	1	4.5 ± 2.2 a	3.3 ± 3.8 ab	82.2 ± 2.2 c	11.1 ± 4.4 a	4.4 ± 4.4 a	84.5 ± 4.4 c

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$. The equally named columns of the two cultivars are estimated together.

Table 2. Factors affecting the culture establishment.

		‘Faassens Black’			‘Drummondii’		
Month of harvesting	PGRs (mg l ⁻¹)	Explants with developed buds (%)	Explants with elongated shoots (%)	Dead explants (%)	Explants with developed buds (%)	Explants with elongated shoots (%)	Dead explants (%)
Hardwood explants							
September	1.0 Kinetin + 0.1 IBA	0.0 ± 0.0 a	0.0 ± 0.0 a	0.0 ± 0.0 a	0.0 ± 0.0 a	0.0 ± 0.0 a	0.0 ± 0.0 a
Softwood explants							
May	PGRs-free	17.8 ± 4.5 bcd	0.0 ± 0.0 a	0.0 ± 0.0 a	31.1 ± 9.7 d	0.0 ± 0.0 a	0.0 ± 0.0 a
	1.0 BAP	13.3 ± 3.8 abc	0.0 ± 0.0 a	0.0 ± 0.0 a	28.9 ± 5.9 cd	0.0 ± 0.0 a	0.0 ± 0.0 a
	1.0 Zeatin	26.6 ± 3.8 cd	0.0 ± 0.0 a	0.0 ± 0.0 a	17.8 ± 8.0 bcd	0.0 ± 0.0 a	0.0 ± 0.0 a
	1.0 Kinetin	2.2 ± 2.2 ab	0.0 ± 0.0 a	0.0 ± 0.0 a	4.4 ± 4.4 ab	0.0 ± 0.0 a	0.0 ± 0.0 a
Buds with removed scales (1) or with scales (2)							
April (1)	PGRs-free	82.2 ± 2.2 f	24.4 ± 8.0 b	0.0 ± 0.0 a	80.0 ± 4.9 f	11.1 ± 4.4 ab	4.5 ± 2.2 ab
November (2)	PGRs-free	55.5 ± 2.2 e	0.0 ± 0.0 a	26.7 ± 3.8 c	71.1 ± 8.0 ef	0.0 ± 0.0 a	13.3 ± 3.8 b

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$. The equally named columns of the two cultivars are estimated together.

Elongation and multiplication of the shoots

On kinetin enriched WPM all explants formed callus. In some of them it was light-green but in a lot of them it was brown and necrotic outside (Fig. 6, 7). On these variants of the medium from 5.6 ± 5.6% to 22.2 ± 14.7% of explants died and the explants with elongated shoots reached from 44.4% to 50.0% (Fig. 8). It was found that 0.5 mg l⁻¹ Kinetin induced maximal percentage of shoot-forming explants (ĐURKOVIĆ, 1996) but statistical difference between both our treatments and cultivars was not found. The concentration of kinetin did not affect the length of the shoots within the cultivar but in comparison with ‘Drummondii’ they were statistically longer in ‘Faassens Black’ on medium with 0.5 mg l⁻¹ kinetin (14.5 ± 3.5 mm). However, axillary shoots formation was not induced in both treatments and cultivars studied (Table 3).



Figure 6. Axillary shoot formation and callus necrosis on explant from *Acer platanoides* ‘Faassens Black’.




Figure 7. Different types of callus formed. A) Pink or red soft callus, easy to be removed, B) Dark brown callus on the surface, C) Rear occasion of light green callus formation.



Figure 8. Shoot elongation from cultivated buds, from both cultivars ("Faassens Black", left and "Drummondii", right) on WPM supplemented with Kinetin.

Table 3. Elongation of the shoots from buds with removed scales after two subcultures.

Kinetin (mg l ⁻¹)	"Faassens Black"			"Drummondii"		
	Explants with elongated shoots (%)	Length of the shoots (mm)	Dead explants (%)	Explants with elongated shoots (%)	Length of the shoots (mm)	Dead explants (%)
0.5	50.0 ± 9.6 a	14.5 ± 3.5 b	11.1 ± 5.6 a	50.0 ± 9.6 a	6.2 ± 0.9 a	11.1 ± 5.6 a
1.0	50.0 ± 19.2 a	9.9 ± 1.9 ab	5.6 ± 5.6 a	44.4 ± 5.6 a	7.6 ± 2.2 a	22.2 ± 14.7 a

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$. The equally named columns of the two cultivars are estimated together.

Effect of the subculture on multiplication rate

In the end of each subculture, reddish and soft callus was formed on the base of explants. However, its inside part was fresh and green (Fig. 7). The duration of subcultivation in our experiment did not provoke statistical significant necrotizing of the explants in "Faassens Black" and significant rate of the elongated initial explant was found after the 10th subculture. These shoots reached the significantly maximal length after the 8th subculture. It was indicated that rejuvenation in mature trees could sometimes be obtained by repeated subcultures (BALESTER et al., 1990; BONGA & VON ADERKAS, 1992; GIOVANNELLI & GIANNINI, 2000) and the proliferation rate could be increased (BALESTER et al., 1990; SANCHEZ et al., 1991). However, the frequency of the induced axillary shoots was low and was noticed on single explants. Also, their length was small and was not affected by the subculture (Fig. 9).

Opposite tendency was found in "Drummondii". Not only single elongated shoots but all of them gradually died and no viable shoots were found after the 8th subculture (Table 4).




Figure 9. Axillary shoot formation A) longer than 5 mm and B) shorter than 5 mm.

Table 4. Effect of the subculture on multiplication rate.

Subculture (day)	“Faassens Black”					“Drummondii”				
	Dead explants (%)	Reaction of the initial explant		Axillary shoots formed		Dead explants (%)	Reaction of the initial explant		Axillary shoots formed	
		Elongated explants (%)	Length of the explant (mm)	Number	Length (mm)		Elongated explants (%)	Length of the explant (mm)	Number	Length (mm)
28	4.2 ± 4.2 a	8.3 ± 4.2 ab	5.0 ± 0.0 a	0.04 ± 0.04 a	5.0 ± 0.0 abc	6.7 ± 6.7 a	0.0 ± 0.0 a	-	-	-
56	4.2 ± 4.2 a	16.7 ± 8.3 bc	7.2 ± 1.6 a	0.25 ± 0.11 ab	8.0 ± 1.6 bc	8.3 ± 8.3 a	8.3 ± 8.3 ab	44.0 ± 0.0 c	-	-
84	0.0 ± 0.0 a	9.3 ± 0.3 ab	22.7 ± 2.8 b	0.19 ± 0.08 ab	7.3 ± 1.2 abc	16.7 ± 8.3 a	0.0 ± 0.0 a	-	-	-
112	0.0 ± 0.0 a	11.1 ± 2.8 ab	39.5 ± 11.0 c	0.23 ± 0.13 ab	8.0 ± 2.2 c	22.0 ± 11.1 a	0.0 ± 0.0 a	-	-	-
140	12.8 ± 2.6 a	25.7 ± 6.8 c	11.6 ± 2.5 ab	0.15 ± 0.10 ab	5.0 ± 0.0 ab	100.0 ± 0.0 b	-	-	-	-
180	14.3 ± 0.0 a	11.9 ± 6.3 ab	9.6 ± 3.0 ab	0.65 ± 0.17 c	6.1 ± 0.4 abc	-	-	-	-	-

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$. The equally named columns of the two cultivars are estimated together.

Factors affecting the cultivation of juvenile genotypes

Effect of the genotype and subculture on the axillary shoot formation

Reddish and soft callus was formed on the base of explants (Fig. 7). The multiplication rate and the length of the axillary shoots were low, were not affected by the genotype, and all explants from genotypes 1 died (Table 5). Also, the subcultures did not affect significantly the number of the induced shoots and they remained smaller than 5 mm (Fig. 9). Single explants formed axillary shoots in genotype 8 and all explants died after the 3rd subculture (Table 6).

Table 5. Effect of the genotype on the multiplication rate and elongation of the axillary shoots.

Genotype	Dead explants (%)	Number of shoots	Length of shoots (mm)
1	33	0.0 ± 0.0 a	-
2	0	1.0 ± 0.3 b	21.9 ± 4.0 a
3	0	0.3 ± 0.3 ab	7.0
4	0	0.5 ± 0.9 ab	12.0 ± 2.0 a
5	0	1.2 ± 0.4 b	19.5 ± 4.4 a
6	33	0.5 ± 0.9 ab	11.5 ± 2.5 a
8	0	1.1 ± 0.2 b	15.1 ± 2.6 a

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$.

Table 6. Effect of the genotype and subculture on multiplication rate of axillary shoots.

Subculture (days)	Genotype					
	2		5		8	
	Dead explants (%)	Number of shoots	Dead explants (%)	Number of shoots	Dead explants (%)	Number of shoots
14	8.7	0.6 ± 0.2 abc	7.7	0.8 ± 0.4 abc	0	-
28	10.0	0.7 ± 0.2 abc	5.0	0.6 ± 0.2 abc	25	0.5 ± 0.5 ac
42	15.6	0.6 ± 0.2 abc	12.9	0.5 ± 0.2 ab	66	-
56	15.0	1.0 ± 0.3 c	11.7	0.2 ± 0.1 a	100	-
70	12.7	0.4 ± 0.1 a	8.8	0.9 ± 0.3 bc	-	-

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$. The equally named columns of the two cultivars are estimated together.

Effect of PGRs on the axillary shoots formation

The combination of BAP and IBA led to the mortality of some explants and it increased in the highest concentration of BAP. Maximal number and length of axillary shoots was induced on the medium with BAP and GA₃. The combination of these PGRs provoked statistically significant additional formation of axillary shoots but their length remained less than 5 mm in the period of cultivation (Table 7).

Table 7. Effect of PGRs on the number and length of the formed axillary shoots after 40 days of cultivation.

PGRs (mg l ⁻¹)	No of the shoots (> 5 mm)	Length of the shoots (> 5 mm)	No of the shoots (< 5 mm)	Dead explants (%)
3 BAP + 0.1 IBA	1.0 ± 0.4 a	7.2 ± 0.7 ab	1.6 ± 0.5 a	45 ± 12 b
1 BAP + 0.1 IBA	1.2 ± 0.2 ab	6.2 ± 0.4 a	3.1 ± 0.6 b	10 ± 7 a
1 BAP + 1 GA ₃	1.9 ± 0.4 b	8.4 ± 0.6 b	4.7 ± 0.6 c	0 ± 0 a
1 2iP	1.7 ± 0.2 ab	7.1 ± 0.5 ab	2.8 ± 0.4 ab	0 ± 0 a

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$.

LINDÉN & RIIKONEN (2006) demonstrated axillary buds formation in Norway maple with 0.01 to 0.1 µM of TDZ. Kinetin in 0.5 mg l⁻¹ was also effective for the induction of axillary shoots (ĐURKOVIĆ, 1996). It was reported that 2 µM BAP promoted higher rate of shoots regeneration and longer shoots in 'Crimson Sentry' in comparison with other cytokinins (LATTIER et al., 2013). The type of cytokinin does not affect the elongation of the initial explant within the genotype. When BAP was used, elongation of the shoots in genotype 2 was not observed and statistical difference in the length of shoots in genotype 5 and "Faassens Black" was not found. Higher number of axillary shoots in genotype 5 was received after the treatment with Kinetin, but different cytokinins do not affect significantly their number in genotype 2 and "Faassens Black". Generally, the quantity of induced axillary shoots was small. The elongation of the shoots was very slow and few of them reached a bit more than 5 mm (Table 8, Fig. 9).

Possibly due to the apical dominance of the initial explant, the formation of axillary shoots and their elongation is limited. Also, different responses to cytokinins could be attributed to the reaction of different genotypes. For this reason the *in vitro* proliferation and elongation of the axillary shoots in *Acer platanoides* and its cultivars remained difficult.

Table 8. Effect of genotype and cytokinins on the reaction of initial explant, number, and length of the formed axillary shoots.

Genotype	PGRs (mg l ⁻¹)	Reaction of the initial explant		Axillary shoots formed	
		Number of elongated explants	Length of the explant (mm)	Number	Length (mm)
Genotype 2	1 BAP	0	-	1.1 ± 0.3 bc	-
	1 2iP	2	12.5 ± 1.6 a	1.1 ± 0.5 bc	-
	2 Kinetin	2	14.6 ± 3.8 a	1.6 ± 0.5 cd	6.0 ± 0.4 a
Genotype 5	1 BAP	4	19.4 ± 5.1 ab	1.1 ± 0.4 bc	8.8 ± 1.3 a
	1 2iP	7	20.5 ± 3.6 ab	0.1 ± 0.1 a	-
	2 Kinetin	2	11.8 ± 1.1 a	2.1 ± 0.6 d	-
'Faassens Black'	1 BAP	6	18.5 ± 3.6 ab	0.0 ± 0.0 a	-
	1 2iP	7	27.4 ± 3.2 b	0.0 ± 0.0 a	-
	2 Kinetin	5	19.4 ± 3.2 ab	0.3 ± 0.2 ab	-

The means (M) ± standard error (SE) within a column followed by the same letter are not significantly different estimated by One-Way ANOVA followed by a post hoc LSD test at $p \leq 0.05$.

ACKNOWLEDGEMENTS

This research was supported by the grant No BG051PO001-3.3.06-0056, financed by the Human Resources Development Operational Programme (2007-2013) and co-financed jointly by the ESF of the EU and the Bulgarian Ministry of Education and Science.

REFERENCES

- ARTJUSHENKO Z. T., VASILEV I. V., GZIRJAN M. S., GOLOVACH A. G., GRUBOV V. I., ZAMJATNIN V. N., PIDOTTI O. A., PILIPENKO F. S., POLETIKO O. M., RODIONENKO G. I., RUSANOV F. N., SAAKOV S. G., SOKOLOV S. Y., FEDOROV A. A., SHULIGINA V. V., SHUHOBODSKII B. A. 1958. Trees and shrubs. *Academy of Science. USSR*: 428-430. [in Russian].
- BALLESTER A., SÁNCHEZ M. C., VIEITEZ A. M. 1990. Development of rejuvenation methods for in vitro establishment, multiplication and rooting of mature trees. In: Rodriguez R., Sánchez-Tamés R., Durzan D. J. (Eds.). *Plant Aging: Basic and Applied Approaches*. Academic Press, New York: 43-49.
- BONGA J. 1982. Tissue culture techniques. In: Bonga J. M., Durzan D. J. (Eds.). *Tissue culture in forestry*. Martinus Nijhoff/Dr. W. Junk Publishers. The Hague: 4-35.
- BONGA J. M. & VON ADERKAS P. 1992. *In vitro culture of trees*. Kluwer Academic Publisher. 236 pp.
- CHENG T.-Y. 1978. Clonal propagation of woody plant species through tissue culture techniques. *Combined Proceedings, International Plant Propagators' Society*. **28**: 139-155.
- CHONG C. & DAIGNEAULT L. 1986. Influence of IBA concentrations on rooting of woody perennial nursery stock. *Combined Proceedings, International Plant Propagators' Society*. **36**: 108-115.
- COMPTON M. E. 1994. Statistical methods suitable for the analysis of plant tissue culture data. *Plant Cell, Tissue and Organ Culture*. **37**: 217-242.
- CONCIOIU M. E., DUTA M., OPREA M. I., TEODORESCU A. 2010. In vitro behavior of 'Globosum' and 'Crimson King' *Acer platanoides* ornamental varieties during initial stage. *Scientific Papers of R.I.F.G.* **26**: 152-156.
- DIRR M. A. 1998. *Acer platanoides* L. Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation and uses. *Fifth Edition*. Stipes Publishing L. L. C. Champaign. Illinois: 40-43.
- DIRR M. A. & HEUSER C. W., JR. 1987. *The reference manual of woody plant propagation: from seed to tissue culture*. Varsity Press. Athens. GA. 239 pp.
- ĐURKOVIĆ J. 1996. In vitro regeneration of Norway maple (*Acer platanoides* L.). *Biologia plantarum*. **38**: 303-307.
- ĐURKOVIĆ J. (2003). Regeneration of *Acer caudatifolium* Hayata plantlets from juvenile explants. *Plant Cell Reports*. **21**: 1060-1064.
- GIONANNELLI A. & GIANNINI R. 2000. Reinvigoration of mature chesnut (*Castanea sativa*) by repeated grafting and micropropagation. *Tree Physiology*. **20**: 1243-1248.
- HARTMANN H. T., KESTER D. E., DAVIES F. T., GENEVE R. L. 2002. *Hartmann and Kester's plant propagation. Principles and practices*. Seventh edition. Prentice Hall. Upper Saddle River, New Jersey.
- HOWARD B. H. 1993. Investigation into inconsistent and low bud-grafting success in *Acer platanoides* 'Crimson King'. *Journal of Horticultural Science*. **68**: 455-462.
- HOWARD B. H., SKENE D. S., COLES J. S. 1974. The effects of different grafting methods upon the development of one-year-old nursery apple trees. *Journal of Horticultural Science*. **49**: 287-295.
- HOWARD B. H. & OAKLEY W. 1997. Bud-grafting in *Acer platanoides* 'Crimson King' related to root growth. *Journal of Horticultural Science*. **72**: 697-704.
- ILIEV I. & BORISOWA D. 2002. Biotechnological methods for improvement and cloning of ornamental species in vitro. *Forestry Ideas*. **3-4**: 104-118 [in Bulgarian].
- ILIEV I., GAJDOSOVA A., LIBIAKOVA G., JAIN S. M. 2010. Plant micropropagation. In: Davey M. and Anthony P. (Eds.). *Plant Cell Culture: Essential Methods*. John Wiley & Sons Ltd.: 1-23.
- KOLESNIKOV A. I. 1958. *Ornamental forms in tree species*. Ministry of municipal economy of RSFSR. Moskow. 272 pp. [in Russian].
- KRÜSSMAN G. 1984. *Acer platanoides* L. In: *Manual of cultivated broad-leaved trees & shrubs*. A-D, Timber Press, Inc. **1**: 92-96.
- LATTIER J. D., TOUCHELL D. H., RANNEY T. G. 2012. Micropropagation of *Acer platanoides* L. 'Crimson Sentry'. SNA research conference. **57**: 296-300.
- LATTIER J. D., TOUCHELL D. H., RANNEY T. G., SMITH J. C. 2013. Micropropagation and polyploid induction of *Acer platanoides* 'Crimson Sentry'. *Journal of Environmental Horticulture*. **31**: 246-252.
- LINDÉN L. & RIIKONEN A. 2006. Effects of 6-benzylaminopurine, thidiazuron and typeof explant on in vitro shoot development of *Acer platanoides* L. *Propagation of Ornamental Plants*. **6**: 201-204.
- LINSMAIER E. M. & SKOOG F. 1965. Organic growth factor requirements of tobacco tissue cultures. *Physiologia Plantarum*. **18**: 100-127.
- LLOYD G. & MCCOWN B. 1980. Commercially feasible micropropagation of mountain laurel (*Kalmia latifolia*) by use of shoot-tip culture. *Combined Proceedings, International Plant Propagators' Society*. **30**: 421-427.
- MARKS T. R. & SIMPSON S. E. 1990. Reduced phenolic oxidation at culture initiation in vitro following the expose of field-grown stockplants to darkness or low levels of irradiance. *Journal of Horticulture Science*. **65**: 103-111.
- MARKS T. R. & SIMPSON S. E. 1994. Factors affecting shoot development in apically dominant *Acer* cultivars in vitro. *Journal of Horticulture Science*. **69**: 543-556.
- PREECE J. E., HUETTEMAN C. A., ASHBY W. C., ROTH P. L. 1991. Micro- and cutting propagation of silver maple. I. Results with adult and juvenile propagules. *Journal of the American Society for Horticultural Science*. **116**: 142-148.

- ROHR R. & HANUS D. 1987. Vegetative propagation of wavy grain sycamore maple. *Canadian Journal of Forest Research.* **17:** 418-420.
- ROVINĂ E. A., CĂLINESCU M., CATIȚA P., ISAC V. 2010. In vitro regeneration capacity of ornamental varieties related to the culture media. *Journal of Horticulture, Forestry and Biotechnology.* **14:** 13-19.
- SANTAMOUR F. S., MCARDLE J. J., MCARDLE A. J. 1982. Checklist of cultivated maples III. *Acer platanoides L.* *Journal of Arboriculture.* **8:** 241-245.
- SÁNCHEZ M. C. & VIEITEZ A. M. 1991. In vitro morphogenetic competence of basal sprouts and crown branches of mature chesnut. *Tree Physiology.* **8:** 59-70.
- ŠEDIVÁ J. 2009. Influence of explant type, sucrose and IBA on in vitro growth of *Acer platanoides L.* 'Jirka'. *Acta Horticulturale.* **812:** 185-188.

Vladimir TOMOV, Ivan ILIEV

Laboratory of Biotechnology and Molecular Genetics,
University of Forestry, 10 Kliment Ohridski blvd., 1797 Sofia, Bulgaria.
E-mail: ivilievltu@yahoo.com

Received: March 26, 2014

Accepted: April 19, 2014