

AMYLASES AS BIOLOGICALLY ACTIVE SUBSTANCES PRODUCED BY BACTERIAL STRAINS COLLECTED FROM POLLUTED AREAS

CÎRSTEA Doina Maria, CIOBOIU Olivia,
ȘTEFĂNESCU Mugur Cristian, CISMAȘIU Carmen Mădălina

Abstract. Amylases are among the most important enzymes, with applications in food, pharmaceutical and textile industries. The present study aimed to render the ability of bacterial strains isolated from environments contaminated with heavy metals or petroleum products to synthesize amylases. Three of the strains studied are able to synthesize amylases; two of them belong to the genus *Pseudomonas* and one was classified in the genus *Bacillus*. Enzymatic activity of these three bacterial strains (*Pseudomonas* sp. HM2, *Pseudomonas* sp. OP16 and *Bacillus* sp. OP3) is favourable and *Bacillus* strain sp. OP3 may represent the object of new applicative studies.

Keywords: biologically active substances, bacteria, amylases.

Rezumat. Amilazele ca substanțe biologice active produse de tulpini bacteriene prelevate din zone poluate. Amilazele sunt printre cele mai importante enzime, cu aplicabilitate în industriile alimentară, farmaceutică și textilă. Prin prezentul studiu s-a urmărit capacitatea unor tulpini bacteriene izolate din medii contaminate cu metale grele sau cu produși petrolieri de a sintetiza amilaze. Trei dintre tulpinile luate în studiu au capacitatea de a sintetiza amilaze, două dintre acestea aparținând genului *Pseudomonas*, iar una fiind afiliată genului *Bacillus*. Activitatea enzimatică, la cele trei tulpini bacteriene (*Pseudomonas* sp. HM2, *Pseudomonas* sp. OP16 și *Bacillus* sp. OP3) este favorabilă, tulpina de *Bacillus* sp. OP3 pretându-se la noi studii cu potențial aplicativ.

Cuvinte cheie: substanțe biologice active, bacterii, amilaze.

INTRODUCTION

Amylases are among the most important enzymes (NAIDU & SARANRAJ, 2013); those amylolytic enzymes are of great significance in food, textile, paper, fermentation, pharmaceutical and sugar industries (KUNAMNENI et al., 2005; WINDISH & MHATRE, 2012).

Amylases can be found in plants, animals, and microorganisms (BEN MASSOUD et al., 1999; OSFAR & TRI, 2011). Amylases can be obtained from: yeast, bacteria, actinomycetes and several fungi (DAMODARA et al., 2012). Among bacterial sources, we can mention: *Bacillus* genus with *B. subtilis*, *B. stearothermophilus*, *B. licheniformis* and *B. amyloliquefaciens*, and some bacteria of the genus *Rhodothermus*, *Lactobacillus* (MONTEIRO DE SOUZA & MAGALHÃES PÉROLA, 2010), *Pseudomonas stutzeri* (MAALEJ et al., 2014).

MATERIAL AND METHODS

The biological material consisted in 25 bacterial strains isolated from areas contaminated with petroleum products and heavy metals.

The selection of the bacterial strains, which have the potential to synthesize amylases, was made on average with starch nutrient agar medium, for 48 hours at 28 °C. After incubation, the plates were flooded with Lugol solution; if starch hydrolysis occurs, a yellow hollow appears around the colonies (LAZĂR et al., 2004).

The bacterial strains were submitted to the protocol for determining the Gram appurtenance, using Gram microbiology technique (LAZĂR et al., 2004); taxonomic identification was made using Biologic GN2 and GP2 micro plates. Micro plates were inoculated at 28 °C for 24 hours, according to the manufacturer's recommendations (Biologic, Inc).

Enzymatic activity was determined after the method described by LAZĂR et al. (2004), using the spectrophotometric method, reading the samples values at 580 nm.

Bacterial cell was cultivated on liquid nutrient agar medium and incubated at 28 °C on a rotary shaker (150 rpm), for five days.

Amylases activity by Zymography was made by gel on a thin 2 % agarose and 1 % soluble starch in Millis-Q water. In the gel were carried holes, which was added 8 µl supernatant. The gel was incubated at 60 °C for 60 minutes and then flooded with Lugol.

RESULTS AND DISCUSSIONS

All bacterial strains were cultivated by starch nutrient agar medium, at 28 °C for 48 hours. After flooding plates with Lugol solution, only 5 bacteria strains gave positive reaction (Fig. 1), which means that 40 % of the strains have the ability to synthesize amylases.




Figure 1. Amylases - synthesis by the bacterial strains.

The five amylolytic bacteria have been identified by using Biologic Systems. Because two bacterial strains were Gram-negative and three Gram-positive, it was used Biologic CN2 micro plates for Gram negative bacteria and CP2 micro plates for Gram positive bacteria. The results based on this biochemical tests indicated that two strains belong to the genus *Pseudomonas* and three were classified in the genus *Bacillus*. It is known that amylases are mainly derived from the genus *Bacillus* (MONTEIRO DE SOUZA & MAGALHÃES PÉROLA, 2010).

After having established the affiliation of the amylolytic bacterial strains, we can note: *Pseudomonas* sp. HM2, *Pseudomonas* sp. OP 16, *Bacillus* sp. OP 3, *Bacillus* sp. OP 9, *Bacillus* sp. OP 6 (indicative: HM - heavy metals, OP - oil pollution and experimental number).

Following the evaluation of enzymatic activity (Table 1), enzymatic activities was identified in three of the five bacterial strains studied at the first screening. All five strains have amylolytic activity.

Table 1. Enzyme activity of amylolytic bacterial strains.

Bacterial strains	UE (enzyme activity unit)				
	24 h	48 h	72 h	96 h	120 h
<i>Pseudomonas</i> HM2	2.514	5.88	4.22	10.02	10.88
<i>Pseudomonas</i> OP16	1.961	6.55	7.352	14.35	11.6
<i>Ps. aeruginosa</i>	0.05	0.21	0.01	0.001	0.09
<i>Bacillus</i> OP 3	2.53	9.099	21.71	24.03	22.01
<i>Bacillus</i> OP 9	0.412	0.3814	0.21	0.21	0.02
<i>Bacillus</i> OP 6	1.141	0.511	0.41	0.43	0.39

The bacterial strains *Pseudomonas* sp. HM2, *Pseudomonas* sp. OP16 have a low enzyme activity compared with the strain *Bacillus* sp. OP3 which has an enzymatic activity similar with other strains of the genus *Bacillus* (OSFAR & TRI, 2011). *Bacillus* sp. OP9 and *Bacillus* sp. OP6 do not show enzymatic activity similar to *Ps. aeruginosa*, a negative control; this bacterial strain cannot metabolize starch.

These results prompted us to test the enzymatic activity of the supernatant by zymography method (Fig. 2).



Figure 2. Amylase activity by Zymography.

The results of amylase activity by zymography of the bacterial strains confirm previous data; *Bacillus* sp. OP 9 and *Bacillus* sp. OP 6 strains do not have enzymatic activity.

CONCLUSIONS

There have been isolated from contaminated environments, three bacterial strains: *Pseudomonas* sp. HM2, *Pseudomonas* sp. OP16 that are dependent on enzymes for metabolising carbohydrates of the starch type. Bacterial strains, *Bacillus* sp. OP3 show a higher enzyme activity compared with the other two bacterial strains (*Pseudomonas* sp. HM2, *Pseudomonas* sp. OP16).

ACKNOWLEDGEMENT

The study was funded by project no. RO1567-IBB05/2015 of the Institute of Biology Bucharest of the Romanian Academy. The authors are grateful to Ana Dinu and Paula Dicul for technical support.

REFERENCES

- BEN MASSOUD E., BEN ALI M., ELLUCH S. 1999. Purification and properties of maltoheptose and maltohexose forming amylase produced by *Bacillus subtilis*. *Enzyme and Microbial Technology*. London. **34**: 662 – 666.
- DAMODARA R. M., RATHNAM B. V. V., PURNIMA A., AYYANNA C. 2012. Affinity chromatography of amylase from *Bacillus licheniformis*. *Enzyme and Microbial Technology*. Springer. Stuttgart. **37**: 712 – 717.
- MAALEJ HANA, AYED HANEN B., GHORBEL-BELLAJ OLFA, MONCEF N., NOOMEN HM. 2014. Production and Biochemical Characterization of a High Maltotetraose (G4) Producing Amylase from *Pseudomonas stutzeri* AS22. *Bio Med Research International*. London: 1-11.
- KUNAMNENI A., PERMAUL K., SINGH S. 2005. Amylase production in solid state fermentation by the thermophilic fungus *Thermomyces funginosus*. *Journal of Bioscience and Bioengineering*. Osaka. **100**(2): 168 - 171.
- LAZĂR V., HERLEA VICTORIA, CERNAT RAMONA, BALOTESCU MARIANA CARMEN, BULAI DOINA, MORARU ANCA. 2004. *Microbiologie generală*. Edit. Universității București. 70-71, 99, 197-198 pp.
- MONTEIRO DE SOUZA PAULA & MAGALHÃES PÉROLA DE OLIVEIRA. 2010. Application of microbial α -amylase in industry – A review. *Brazilian Journal of Microbiology*. Sao Paulo. **41**(4): 850–861.
- NAIDU M. A. & SARANRAJ P. 2013. Bacterial Amylase: A Review. *International Journal of Pharmaceutical & Biological Archives*. London. **4**(2): 274 – 287.
- OSFAR SJOFJAN & TRI ARDYATI. 2011. Extracellular Amylase Activity of Amylolytic Bacteria Isolated from Quail's (*Coturnix japonica*) Intestinal Tract in Corn Flour Medium. *International Journal of Poultry Science*. London. **10**(5): 411-415.
- WINDISH W. W & MHATRE N. S. 2012. Microbial amylases. In: Wayne WU, (edit.). *Advances in applied microbiology*. Academic Press. New York. **7**: 273 - 304.

Cîrstea Doina Maria, Ștefănescu Mugur Cristian, Cismașiu Carmen Mădălina

Institute of Biology Bucharest of the Romanian Academy,
296 Splaiul Independenței, 060031, Bucharest, Romania.

E-mail: maria_2na@yahoo.com

Cioboiu Olivia

The Oltenia Museum Craiova, Str. Popa Șapcă No. 8, 200422, Craiova, Romania.

E-mail: oliviacioboiu@gmail.com; cioboiu.olivia@yahoo.com

Received: March 31, 2015

Accepted: June 23, 2015